




PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
 International Bureau


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 451/02, A01N 43/40		A1	(11) International Publication Number: WO 96/37494
			(43) International Publication Date: 28 November 1996 (28.11.96)
(21) International Application Number: PCT/GB96/01151		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 13 May 1996 (13.05.96)			
(30) Priority Data: 9510459.2 24 May 1995 (24.05.95) GB			
(71) Applicant (for all designated States except US): ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB).			
(72) Inventors; and		Published	
(75) Inventors/Applicants (for US only): URCH, Christopher, John [GB/GB]; 40 Dalcross, Crown Wood, Bracknell, Berkshire RG12 3UJ (GB). SALMON, Roger [GB/GB]; 38 Tawfield, Bracknell, Berkshire RG12 4YU (GB). LEWIS, Terence [GB/GB]; 18 Quintilis, Roman Hill, Bracknell, Berkshire RG12 4QQ (GB). GODFREY, Christopher, Richard, Ayles [GB/GB]; 159 Viking, Great Hollands, Bracknell, Berkshire RG12 8UW (GB). CLOUGH, Martin, Stephen [GB/GB]; 25 Stephen Close, Twyford, Berkshire RG10 0XN (GB).		With international search report.	
(74) Agents: BISHOP, Nigel, Douglas et al.; Zeneca Agrochemicals, Intellectual Property Dept., Jealott's Hill Research Station, P.O. Box 3538, Bracknell, Berkshire RG42 6YA (GB).			

(54) Title: **BICYCLIC AMINES AS INSECTICIDES**



(I)



(A)

(57) Abstract

A compound of formula (I), wherein R^1 represents a group of formula (A) where each of W, X, Y and Z represents either a group CR or the nitrogen atom, provided that not more than two of W, X, Y and Z represent the nitrogen atom and where each R present is independently selected from hydrogen and halogen atoms and cyano, amino, hydrazino, acylamino, hydroxy, alkyl, hydroxyalkyl, alkoxy, haloalkyl, haloalkoxy, alkenyl, alkenyloxy, alkoxyalkenyl, alkynyl, carboxylic acyl, alkoxycarbonyl, aryl and heterocyclyl groups, said groups comprising up to 6 carbon atoms, and wherein R^2 represents hydrogen or cyano or a group selected from alkyl, aryl, heteroaryl, aralkyl, heteroarylalkyl, alkenyl, aralkenyl, alkynyl, alkoxycarbonyl, alkanesulfonyl, arenesulfonyl, alkanyloxy, carbonyl, aryloxy, heterocyclylalkyl, carbamyl or dithiocarboxyl groups, said groups comprising from 1 to 15 carbon atoms, said groups being optionally substituted with one or more substituents selected from, halogen, cyano, carboxyl, carboxylic acyl, carbamyl, alkoxycarbonyl, alkoxy, alkylenedioxy, hydroxy, nitro, haloalkyl, alkyl, amino, acylamino, imidate and phosphonate groups; and acid addition salts and quaternary ammonium salts and N-oxide derived therefrom. The compounds are useful as insecticides.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

BICYCLIC AMINES AS INSECTICIDES

This invention relates to novel bicyclic amines, to processes for preparing them, to insecticidal compositions comprising and to methods of combating and controlling insect pests therewith.

- 5 The invention provides compounds of formula (I) wherein R^1 represents a group of formula (A) where each of W, X, Y and Z and Z represents either a group CR or the nitrogen atom, provided that not more than two of W, X, Y and Z represent the nitrogen atom and where each R present is independently selected from hydrogen and halogen atoms and cyano, amino, hydrazino, acylamino, hydroxy, alkyl, hydroxyalkyl, alkoxy, haloalkyl, haloalkoxy, alkenyl, alkenyloxy, alkoxyalkenyl, alkynyl, carboxylic acyl, alkoxycarbonyl, aryl and heterocyclyl groups, said groups comprising up to 6 carbon atoms, and wherein R^2 represents hydrogen or cyano or a group selected from alkyl, aryl, heteroaryl, aralkyl, heteroarylalkyl, alkenyl, aralkenyl, alkynyl, alkoxycarbonyl, alkanesulfonyl, arenesulfonyl, alkanyloxycarbonyl, aralkyloxycarbonyl, aryloxycarbonyl, heterocyclylalkyl, carbamyl or dithiocarboxyl groups, said groups comprising from 1 to 15 carbon atoms, said groups being optionally substituted with one or more substituents selected from, halogen, cyano, carboxyl, carboxylic acyl, carbamyl, alkoxycarbonyl, alkoxy, alkylenedioxy, hydroxy, nitro, haloalkyl, alkyl, amino, acylamino, imidate and phosphonato groups; and acid addition salts and quaternary ammonium salts and N-oxides derived therefrom. R^1 is preferably a halo-
- 15 substituted phenyl, pyridyl or diazinyl group.
- 20

In a preferred aspect invention provides compounds of formula (I) where R^1 represents an optionally halogen substituted phenyl group or an optionally halogen substituted pyridyl, pyridazinyl or pyrazinyl group and R^2 represents hydrogen or a C_{1-6} alkyl, alkenyl, alkynyl, phenyl, benzyl, pyridylmethyl, thienylmethyl, thiazolylmethyl group which may be optionally substituted with one or more alkyl, alkoxy, alkoxycarbonyl, cyano, optionally substituted alkane sulphonyl groups or halogen atoms; and acid addition salts thereof.

25

One particularly preferred group of compounds are those wherein R^1 represents an optionally halogen substituted phenyl or pyridyl group and R^2 represents a alkyl group containing up to 4 carbon atoms which may optionally be substituted with one or more halogen atoms.

30

An especially preferred group of compounds are those wherein R¹ represents a 5-halopyrid-3-yl group and R² represents hydrogen or a haloalkyl, haloalkenyl or halobenzyl group.

Specific compounds of formula I according to the invention include those set out in Table I below in which the groups represented by R¹ and R² are given for each compound, together with the melting point (°C) or an indication of the physical state of the compound.

TABLE I

Compound No	R ¹	R ²	Melting Point
1	3,5-dichlorophenyl	methyl	145-146 °C
2	3,5-difluorophenyl	methyl	93-94 °C
3	2,3-difluorophenyl	methyl	oil
4	pentafluorophenyl	methyl	oil
5	2,3-dichlorophenyl	methyl	solid
6	4-methoxyphenyl	benzyl (Isomer A)	178.4 °C
7	4-methoxyphenyl	benzyl (Isomer B)	95-100 °C
8	phenyl	benzyl	90-90.5 °C
9	3,5-difluorophenyl	H	112.1 °C
10	3,5-difluorophenyl	benzyl	85.2 °C
11	3,5-difluorophenyl	5,6-dichloropyrid-3-ylmethyl	143.2-144.2 °C
12	3,5-difluorophenyl	pyrid-2-ylmethyl	127.9-128.5 °C
13	3,5-difluorophenyl	3-methylbenzyl	95.9-96.1 °C
14	3,5-difluorophenyl	4-chlorobenzyl	95.5-96.7 °C
15	3,5-difluorophenyl	pyrid-3-ylmethyl	78.2 °C
16	3,5-difluorophenyl	3,4-methylenedioxybenzyl	oil
17	3,5-difluorophenyl	3,5-dichlorobenzyl	154.1 °C
18	3,5-difluorophenyl	3,3-difluoroprop-2-en-1-yl	94.6 °C
19	3,5-difluorophenyl	2-hydroxy-2-phenylethyl	120.8 °C
20	3,5-difluorophenyl	1-phenyl-2-hydroxyethyl	168.8 °C
21	3,5-difluorophenyl	allyl	70.5 °C
22	3,5-difluorophenyl	propargyl	108.4 °C
23	3,5-difluorophenyl	2-fluoroethyl	oil
24	3,5-difluorophenyl	2-hydroxyethyl	100.4 °C
25	3,5-difluorophenyl	2-methoxyethyl	54.8 °C
26	3,5-difluorophenyl	2-cyanoethyl	115 °C
27	3,5-difluorophenyl	5-chlorothien-2-ylmethyl	114 °C
28	3,5-difluorophenyl	6-chloropyrid-2-yl	gum
29	3,5-difluorophenyl	2-methylthiazol-5-ylmethyl	140 °C
30	3,5-difluorophenyl	2-iminyl-2-methoxyethyl	109 °C
31	phenyl	benzyl (exo-isomer)	115-116 °C
32	phenyl	benzyl (endo-isomer)	97 °C
33	pyrid-3-yl	methyl	87 °C

- 3 -

Compound No	R ¹	R ²	Melting Point
34	pyrid-3-yl	2-fluoroethyl	86-88 °C
35	pyrid-3-yl	allyl	90-92 °C
36	pyrid-3-yl	H	80-81 °C
37	pyrid-3-yl	benzyl	119-120 °C
38	pyrid-3-yl	ethyl	oil
39	pyrid-3-yl	t-butoxycarbonylmethyl	gum
40	N-methylpyridinium-3-yl	t-butoxycarbonyl (iodide)	185-187 °C
41	6-chloropyridazin-3-yl	methyl	119-120 °C
42	pyrid-3-yl	propyl	oil
43	6-chloropyrazin-2-yl	methyl	80 °C
44	pyrid-3-yl	methane-sulphonylmethylsulphonyl	163-164 °C
45	pyrid-3-yl	methane-sulphonyl	135 °C
46	6-chloropyrid-3-yl	methyl	gum
47	pyrid-3-yl	methoxymethyl	oil
48	pyrid-3-yl	ethoxymethyl	oil
49	pyrid-3-yl	cyanomethyl	90-91 °C
50	pyrid-3-yl	ethoxycarbonylmethyl	gum
51	pyrid-3-yl	methoxycarbonylmethyl	gum
52	2-fluoro-4-nitrophenyl	methyl	100-102 °C
53	3-fluorophenyl	methyl	oil
54	pyrid-3-yl	2-hydroxyethyl	155.2-156.8 °C
55	5,6-dichloropyrid-3-yl	methyl	110.1-111.4 °C
56	pyrid-3-yl	propargyl	119-8-121.1 °C
57	pyrid-3-yl	methyl	gum
58	pyrid-3-yl	but-2-en-1-yl	193-194 °C
59	3,5-difluorophenyl	4-nitrophenyl	96.9-97.9 °C
60	5-chloropyrid-3-yl	methyl	152.8-154.5 °C
61	pyrid-3-yl	phenyl	136-137 °C
62	pyrazin-2-yl	methyl	76-76.9 °C
63	2,6-dichloropyrimid-4-yl	methyl	95.3-96.8 °C
64	5-chloropyrid-3-yl	2-fluoroethyl	125.9-126.9 °C
65	2,6-dichloropyrid-4-yl	methyl	165-165.8 °C
66	2-chloro-6-hydrazinopyrid-4-yl	methyl	72-73 °C
67	pyrid-4-yl	methyl	74.5-76.1 °C
68	5-bromopyrid-3-yl	methyl	144.1-145.2 °C
69	5-chloropyrid-3-yl	vinylloxycarbonyl	gum
70	5-chloropyrid-3-yl	H	85-87 °C
71	6-chloropyrid-2-yl	methyl	103.9-104.8 °C
72	5-chloropyrid-3-yl	2,2,2-trifluoroethyl	109.5-111.5 °C
73	3,5-difluorophenyl	pyrid-2-yl	oil
74	5-chloropyrid-3-yl	phenyl	122-123 °C
75	5-chloropyrid-3-yl	propargyl	110-112 °C
76	5-chloropyrid-3-yl	allyl	78-80 °C

- 4 -

Compound No	R ¹	R ²	Melting Point
77	5-methoxypyrid-3-yl	methyl	112.2-113.1 °C
78	5-chloropyrid-3-yl	ethyl	116-118 °C
79	5-chloropyrid-3-yl	butyl	48-50 °C
80	5-ethoxypyrid-3-yl	methyl	567.2-57 °C
81	5-chloropyrid-3-yl	hexyl	resin
82	5-chloropyrid-3-yl	phenoxycarbonyl	117-123 °C
83	5-chloropyrid-3-yl	2,2,2-trichloroethoxycarbonyl	oil
84	5-chloropyrid-3-yl	ethoxycarbonyl	oil
85	5-chloropyrid-3-yl	fluoren-9-ylmethyloxycarbonyl	68-70 °C
86	5-chloropyrid-3-yl	ethoxycarbonylmethyl	gum
87	5-chloropyrid-3-yl	isopropyl	oil
88	5-chloropyrid-3-yl	4,4,4-trifluorobut-3-on-1-en-1-yl	143.9-145.1 °C
89	5-chloropyrid-3-yl	1-methyl-2,2,2-trichloroethoxycarbonyl	152-155 °C
90	5-chloropyrid-3-yl	allyloxycarbonyl	oil
91	5-chloropyrid-3-yl	benzyloxycarbonyl	oil
92	5-chloropyrid-3-yl	2-chloroethoxycarbonyl	gum
93	5-chloropyrid-3-yl	pentafluorobenzyl	143-144 °C
94	5-chloropyrid-3-yl	4-nitrophenyl	213-214.5 °C
95	5-chloropyrid-3-yl	acetyl	162-165 °C
96	5-chloropyrid-3-yl	trifluoroacetyl	121-124 °C
97	5-chloropyrid-3-yl	4-chlorobenzoyl	175-177 °C
98	5-chloropyrid-3-yl	4-fluorobenzoyl	200-204 °C
99	5-chloropyrid-3-yl	3-fluoropropyl	oil
100	5-chloropyrid-3-yl	2,4-bis(trifluoromethyl)benzyl	112-114 °C
101	5-chloropyrid-3-yl	4-carboxybenzyl	gum
102	5-(prop-1-enyloxy)pyrid-3-yl	methyl	gum
103	5-chloropyrid-3-yl	2,3-difluorobenzyl	102-103 °C
104	5-chloropyrid-3-yl	2-phenylethyl	oil
105	5-chloropyrid-3-yl	4-cyanophenyl	201-204 °C
106	5-chloropyrid-3-yl	3,3-difluoroprop-2-en-1-yl	oil
107	5-chloropyrid-3-yl	carboxymethyl	165-167 °C
108	5-chloropyrid-3-yl	3,5-dibromobenzyl	194-196 °C
109	5-chloropyrid-3-yl	3-chloro-4-fluorobenzyl	95-97 °C
110	5-chloropyrid-3-yl	formyl	141-142 °C
111	5-chloropyrid-3-yl	isopropoxycarbonyl	gum
112	5-chloropyrid-3-yl	benzenesulfonyl	210-211 °C
113	5-chloropyrid-3-yl	2,4,6-trifluorobenzyl	106-107 °C
114	5-chloropyrid-3-yl	2,3,6-trifluorobenzyl	125-127 °C
115	5-chloropyrid-3-yl	1-cyano-1-phenylmethyl	141-142 °C
116	5-chloropyrid-3-yl	methoxycarbonyl	oil

- 5 -

Compound No	R ¹	R ²	Melting Point
117	5-chloropyrid-3-yl	pyrid-2-ylmethyl	123-125 °C
118	5-chloropyrid-3-yl	pyrid-3-ylmethyl	105-107 °C
119	5-chloropyrid-3-yl	pyrid-4-ylmethyl	111-114 °C
120	pyrid-2-yl	2-fluoroethyl	82-84 °C
121	5-chloropyrid-3-yl	(R)-1-phenylethyl	115.6-116.7 °C
122	5-chloropyrid-3-yl	(S)-1-phenylethyl	113.4-115 °C
123	5-chloropyrid-3-yl	2-methylthiazol-4-ylmethyl	81-83 °C
124	5-chloropyrid-3-yl	3,5-dimethylisoxazol-4-ylmethyl	95-99 °C
125	5-chloropyrid-3-yl	5-chlorothien-2-ylmethyl	119-121 °C
126	5-chloropyrid-3-yl	5-trifluoromethylpyrid-2-yl	124.5-125.5 °C
127	pyrid-3-yl	2-methoxyethyl	251-253 °C
128	5-chloropyrid-3-yl	6-fluoropyrid-2-yl	131.5-132.5 °C
129	5-chloropyrid-3-yl	4-fluorophenyl	solid
130	5-chloropyrid-3-yl	2,2,3,3,3-pentafluoropropyl	oil
131	5-chloropyrid-3-yl	2,2,3,3-tetrafluoropropyl	110-113 °C
132	5-chloropyrid-3-yl	2,2,3,3,4,4,4-heptafluoropropyl	oil
133	5-chloropyrid-3-yl	2,2,3,3,4,4,5,5-octafluoropropyl	oil
134	5-aminopyrid-3-yl	methyl	188-190 °C
135	5-chloropyrid-3-yl	1-phenyl-1-carboxamidomethyl	193-195 °C
136	5-chloropyrid-3-yl	6-trifluoromethylpyrid-2-yl	117.5-118.5 °C
137	5-chloropyrid-3-yl	6-chloropyrid-2-yl	176-177 °C
138	5-chloropyrid-3-yl	mercaptothiocarbonyl	224 °C
139	5-chloropyrid-3-yl	t-butyl	127-129 °C
140	5-chloropyrid-3-yl	2-(ethoxycarbonyl)ethyl	gum
141	5-chloropyrid-3-yl	2-carboxyethyl	180-181 °C
142	5-chloropyrid-3-yl	2,2-difluoroethyl	101-104 °C
143	5-bromopyrid-3-yl	2,2,2-trifluoroethyl	105-110 °C
144	5-chloropyrid-3-yl	fluorocarbonyl	165-167 °C
145	5-chloropyrid-3-yl	N-methyl-N-phenyl carbamyl	108-110 °C
146	5-chloropyrid-3-yl	N-t-butylcarbamyl	62-65 °C
147	5-iodopyrid-3-yl	methyl	144-145 °C
148	5-hydroxypyrid-3-yl	methyl	170.9-171.7 °C
149	5-chloropyrid-3-yl	4-morpholinocarbonyl	143-145 °C
150	5-chloropyrid-3-yl	N,N-diisopropylcarbamyl	118-121 °C
151	5-chloropyrid-3-yl	pentafluorophenyl	gum
152	5-chloropyrid-3-yl	6-chloropyrimin-4-yl	174-176 °C
153	5-chloropyrid-3-yl	2-acetamidothiazol-4-ylmethyl	solid
154	5-chloropyrid-3-yl	N-(3-chloro-4-fluorophenyl) carbamyl	216-218 °C
155	5-chloropyrid-3-yl	5-chloropyrid-3-yl	gum

Compound No	R ¹	R ²	Melting Point
156	5-chloropyrid-3-yl	4-trifluoromethylpyrid-3-carboxamidomethyl	149.3-150.4 °C
157	5-chloropyrid-3-yl	4-trifluoromethylpyrid-3-ylcarbonyl	80.3-81.9 °C
158	5-chloropyrid-3-yl	5-chloro-1,2,3-triadiazol-4-ylmethyl	oil
159	5-chloropyrid-3-yl	1-formyl-1-phenylethyl	124-126 °C
160	5-chloropyrid-3-yl	4,4,4-trifluorobutyl	oil
161	5-methoxypyrid-3-yl	2,2,2-trifluoroethyl	88-90 °C
162	5-chloropyrid-3-yl	4-ethoxycarbonylphenyl	131.5-132.5 °C
163	5-chloro-6-fluoro pyrid-3-yl	2,2,2-trifluoroethyl	121-122 °C
164	5-chloropyrid-3-yl	vinylloxycarbonyl	gum
165	5-acetamidopyrid-3-yl	methyl	195-197 °C
166	5-methoxypyrid-3-yl	cyanomethyl	solid
167	5-chloropyrid-3-yl	3-chloromethyl-1,2,4-thiadiazol-5-yl	gum
168	5-chloropyrid-3-yl	5-chlorothiazol-2-yl	111-112 °C
169	5-chloropyrid-3-yl	cyano	168-170 °C
170	5-chloropyrid-3-yl	4-carboxyphenyl	solid
171	5-methoxypyrid-3-yl	vinylloxycarbonyl	gum
172	5-methoxypyrid-3-yl	H	112-114 °C
173	5-chloropyrid-3-yl	4-chlorophenyl	137.5-138 °C
174	5-trifluoromethylpyrid-3-yl	methyl	118.2-118.5 °C
175	5-chloropyrid-3-yl	2-phenylbut-3-en-2-yl	gum
176	5-chloropyrid-3-yl	3-hydroxy-2-phenylprop-2-yl	124-126 °C
177	5-trifluoromethylpyrid-3-yl	formyl	117-121 °C
178	5-chloropyrid-3-yl	3-acetoxy-2-phenylprop-2-yl	130-131 °C
179	5-chloropyrid-3-yl	2-fluoro-2-phenylprop-1-yl	gum
180	5-chloropyrid-3-yl	3,3,5-trimethylhexyl	
181	5-bromopyrid-3-yl	vinylloxycarbonyl	63-66 °C
182	5-chloropyrid-3-yl	pyrimid-2-yl;	148.5-149.5 °C
183	5-trifluoromethylpyrid-3-yl	vinylloxycarbonyl	resin
184	5-trifluoromethylpyrid-3-yl	H	resin
185	pyrid-3-yl	vinylloxycarbonyl	gum
186	5-trifluoromethylpyrid-3-yl	3-chlorobenzyl	oil
187	5-chloropyrid-3-yl	2-chloropyrimid-4-yl	210-212 °C
188	5-chloropyrid-3-yl	4-trifluoromethylphenyl	131-132 °C
189	5-(pyrrol-1-yl)pyrid-3-yl	methyl	gum

Compound No	R ¹	R ²	Melting Point
190	N-oxidopyrid-3-yl	t-butoxycarbonyl	55-57 °C
191	5-chloropyrid-3-yl	2-phenyl-2-isopropylaminoprop-1-yl	gum
192	5-chloropyrid-3-yl	2-phenyl-3-hydroxy-3-cyanoprop-2-yl	172-175 °C
193	5-ethynylpyrid-3-yl	methyl	solid
194	pyrimid-4-yl	methyl	solid
195	5-(1-ethoxyvinyl)pyrid-3-yl	methyl	gum
196	pyrid-3-yl	1,1-dimethylpropyl	gum
197	5-chloropyrid-3-yl	1-ethoxycarbonylethyl	gum
198	5-bromopyrimid-4-yl	methyl	141-145 °C
199	5-trifluoromethylpyrid-3-yl	2,2,2-trifluoroethyl	gum
200	6-pyrimid-4-ylpyrimid-4-yl	methyl	136-154 °C
201	5-acetylpyrid-3-yl	methyl	gum
202	5-fluoropyrid-3-yl	methyl	135-137 °C
203	5-bromopyrid-3-yl	H	128-130 °C
204	5-bromopyrid-3-yl	2-chlorobenzyl	109-111 °C
205	5-chloropyrid-3-yl	2-(3-chlorophenyl)prop-2-yl	gum
206	5-(2-hydroxyprop-2-yl)pyrid-3-yl	methyl	gum
207	5-chloropyrid-3-yl	2-methylbut-3-yn-2-yl	107-110 °C
208	5-bromopyrid-3-yl	ethoxycarbonyl	92-94 °C
209	5-chloropyrid-3-yl	2-methyl-1,1,1-trifluoroprop-2-yl	97-99 °C
210	5-bromopyrid-3-yl	2-methylpropyl	oil
211	5-chloropyrid-3-yl	1-methoxycarbonylethyl (Isomer A)	gum
212	5-chloropyrid-3-yl	1-methoxycarbonylethyl (racemate)	105-106 °C
213	5-chloropyrid-3-yl	1-methoxycarbonylethyl (Isomer B)	gum
214	6-methoxypyrazin-2-yl	methyl	gum
215	5-chloropyrid-3-yl	1-cyano-1-(3-chlorophenyl)methyl	foam
216	5-chloropyrid-3-yl	1-cyanoethyl	gum
217	5-phenylpyrid-3-yl	vinylloxycarbonyl	gum
218	5-chloropyrid-3-yl	4,4-difluorobut-3-en-1-yl	oil
219	5-chloropyrid-3-yl	1-cyano-2-methylprop-1-yl	gum
220	5-phenylpyrid-3-yl	H	gum
221	5-methylpyrid-3-yl	vinylloxycarbonyl	gum
222	5-ethoxycarbonylpyrid-3-yl	vinylloxycarbonyl	solid
223	5-chloropyrid-3-yl	2-cyanoprop-2-yl	solid

- 8 -

Compound No	R ¹	R ²	Melting Point
224	6-ethynylpyrazin-2-yl	methyl	solid
225	5-ethoxycarbonylpyrid-3-yl	H	gum
226	5-(2,2,2-trifluoroethoxy)pyrid-3-yl	trifluoroethyl	oil
227	5-chloropyrid-3-yl	3,5-bis(trifluoromethyl)benzyl	
228	5-chloropyrid-3-yl	2,6-difluorobenzyl	
229	5-chloropyrid-3-yl	3-phenoxybenzyl	
230	5-chloropyrid-3-yl	3-bromo-4-fluorobenzyl	
231	5-chloropyrid-3-yl	3-benzoylbenzyl	
232	5-chloropyrid-3-yl	3-(2,6-dichlorobenzoyl)benzyl	
233	5-chloropyrid-3-yl	3-(2,6-difluorobenzoyl)benzyl	
234	5-chloropyrid-3-yl	4-allyl-2,3,5,6-tetrafluorobenzyl	
235	5-chloropyrid-3-yl	3-trifluoromethoxybenzyl	
236	5-chloropyrid-3-yl	naphth-1-ylmethyl	
237	5-chloropyrid-3-yl	benzyl	
238	5-chloropyrid-3-yl	2-bromobenzyl	
239	5-chloropyrid-3-yl	2-methylbenzyl	
240	5-chloropyrid-3-yl	3-bromobenzyl	
241	5-chloropyrid-3-yl	3-methoxycarbonylbenzyl	
242	5-chloropyrid-3-yl	3-methylbenzyl	
243	5-chloropyrid-3-yl	4-bromobenzyl	
244	5-chloropyrid-3-yl	4-methoxycarbonylbenzyl	
245	5-chloropyrid-3-yl	4- <i>t</i> -butylcarbonylbenzyl	
246	5-chloropyrid-3-yl	4- <i>t</i> -butylbenzyl	
247	5-chloropyrid-3-yl	4-isopropylbenzyl	
248	5-chloropyrid-3-yl	4-methylbenzyl	
249	5-chloropyrid-3-yl	3,4-difluorobenzyl	
250	5-chloropyrid-3-yl	2-fluorobenzyl	
251	5-chloropyrid-3-yl	3-bromo-5-fluorobenzyl	
252	5-chloropyrid-3-yl	2,4-difluorobenzyl	
253	5-chloropyrid-3-yl	3-fluorobenzyl	
254	5-chloropyrid-3-yl	4-fluorobenzyl	
255	5-chloropyrid-3-yl	3-trifluoromethylbenzyl	
256	5-chloropyrid-3-yl	4-trifluoromethylbenzyl	
257	5-chloropyrid-3-yl	2-fluoro-3-chlorobenzyl	
258	5-chloropyrid-3-yl	2-chloro-3,6-difluorobenzyl	
259	5-chloropyrid-3-yl	2-chlorobenzyl	
260	5-chloropyrid-3-yl	2,6-dichlorobenzyl	
261	5-chloropyrid-3-yl	3-chlorobenzyl	
262	5-chloropyrid-3-yl	2-iodo-4-fluorobenzyl	108-109 °C
263	5-chloropyrid-3-yl	2-fluoro-3-methylbenzyl	
264	5-chloropyrid-3-yl	2-(<i>N</i> -succinimido)benzyl	

- 9 -

Compound No	R ¹	R ²	Melting Point
265	5-chloropyrid-3-yl	2-fluoro-5-trifluoromethylbenzyl	
266	5-chloropyrid-3-yl	biphenyl-2-ylmethyl	
267	5-chloropyrid-3-yl	2-cyanobenzyl	
268	5-chloropyrid-3-yl	4-(1,2,3-thiadiazol-4-yl)benzyl	
269	5-chloropyrid-3-yl	3-(4-fluorophenoxy)benzyl	
270	5-chloropyrid-3-yl	4-cyanobenzyl	
271	5-chloropyrid-3-yl	2,3,4-trifluorobenzyl	
272	5-chloropyrid-3-yl	2-nitrobenzyl	
273	5-chloropyrid-3-yl	2-nitro-6-fluorobenzyl	
274	5-chloropyrid-3-yl	3-nitrobenzyl	
275	5-chloropyrid-3-yl	4-nitrobenzyl	
276	5-chloropyrid-3-yl	2-methylprop-1-yl	
277	5-chloropyrid-3-yl	decyl	
278	5-chloropyrid-3-yl	2-phenoxyethyl	
279	5-chloropyrid-3-yl	2-ethoxyethyl	
280	5-chloropyrid-3-yl	3-methylbut-1-yl	
281	5-chloropyrid-3-yl	3-methoxycarbonylprop-1-yl	
282	5-chloropyrid-3-yl	3-phenylprop-1-yl	
283	5-chloropyrid-3-yl	cyclohexylmethyl	
284	5-chloropyrid-3-yl	2-cyanoethyl	
285	5-chloropyrid-3-yl	3-cyanoprop-1-yl	
286	5-chloropyrid-3-yl	2-hydroxyprop-1-yl	
287	5-chloropyrid-3-yl	2-propenoyloxyethyl	
288	5-chloropyrid-3-yl	2-methoxyethyl	
289	5-chloropyrid-3-yl	tetrahydropyran-2-ylmethyl	
290	5-chloropyrid-3-yl	2-hydroxymethylprop-1-yl	
291	5-chloropyrid-3-yl	diethylphosphonomethyl	69-70 °C
292	5-chloropyrid-3-yl	phosphonomethyl	242-245 °C
293	5-chloropyrid-3-yl	methyl (N-oxide)	153-155 °C
294	pyrid-3-yl	t-butoxycarbonyl	
295	6-chloropyrid-3-yl	H	
296	5-chloropyrid-3-yl	methoxy	
297	2-chloropyrimid-4-yl	2,2,2-trifluoroethyl	
298	6-chloropyrazin-2-yl	vinylloxycarbonyl	
299	6-chloropyrazin-2-yl	H	
300	6-chloropyrazin-2-yl	3-chlorobenzyl	
301	6-chloropyrazin-2-yl	cyanomethyl	
302	5-chloropyrid-3-yl	1-(3-chlorophenyl)ethyl	
303	5-chloropyrid-3-yl	4-methoxyphenyl	
304	5-cyanopyrid-3-yl	methyl	
305	2-chloropyrid-4-yl	methyl	
306	5-chloropyrid-3-yl	2-phenylprop-1-en-1-yl	
307	5-chloropyrid-3-yl	methylmercaptothio-carbonyl	

- 10 -

Compound No	R ¹	R ²	Melting Point
308	5-(2,2,2-trifluoroethoxy)pyrid-3-yl	methyl	
309	5-iodopyrid-3-yl	vinylloxycarbonyl	

It will be appreciated that the bicyclic amine compounds of formula I are capable of existing in more than one isomeric form since the groups R¹ and R² may be positioned in either an exo or endo relationship, and the present invention embraces within its scope both

5 exo and endo forms and mixtures thereof and also any further isomeric variants arising from cis and trans substitution patterns or chiral centres present in either of R¹ or R².

Suitable acid addition salts include those with an inorganic acid such as hydrochloric, hydrobromic, sulfuric, nitric and phosphoric acids, or an organic carboxylic acid such as oxalic, tartaric, lactic, butyric, tolúic, hexanoic and phthalic acids, or sulphonic acids such as

10 methane, benzene and toluene sulphonic acids. Examples of salts of compound 72 (Table I) with some less common acids are given in Table IA.

TABLE IA

Compound No	Acid Component
310	2-chlorobenzoic acid
311	4-chlorophenoxyacetic acid
312	2,4,6-trimethylbenzoic acid
313	3-benzylbenzoic
314	4-hydroxybenzoic acid
315	1-phenylpropionic acid
316	3-(4-hydroxyphenyl)propenoic acid
317	undecanoic acid
318	4-(4-hydroxyphenyl)butyric acid
319	2-hydroxy-5-nitrobenzoic acid
320	2-nitro-5-N-methylformamidobenzoic acid
321	2,2,3,3-tetramethylcyclopropanoic acid

15 The preparation of the compounds of formula (I) may be accomplished by use of one or more of the following synthetic techniques described below and further illustrated in the Examples.

The compounds of general formula (I) can be prepared from compounds of general formula (II) by treating them with a suitable base, such as potassium carbonate, in the presence of

20 compound of formula R² where L is a suitable leaving group such as a halide or triflate.

Alternatively, compounds of general formula (I) can be prepared from compounds of general formula (II) by reductive amination with an aldehyde (R^3CHO ; where $R^3CH_2=R^2$) in the presence of a suitable reducing agent such as formic acid.

Compounds of general formula (II) can be prepared by demethylating compound of general formula (III) by, for instance, treating them first with a chloroformate ester (such as vinyl chloroformate) to produce a carbamate, followed by acid hydrolysis.

Compounds of general formula (III) can be prepared by treating 3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (IV) first with a suitable base, such as lithium diisopropylamide (LDA), followed by reaction with an aryl or heteroaryl halide (R^1Hal).

3-Cyano-8-methyl-8-azabicyclo[3.2.1]octane (IV) can be prepared by treating tropinone (V) with tosylmethyl isocyanide in the presence of a suitable base, such as potassium ethoxide. As an alternative 3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (IV) can be prepared from tropine (XII) by treatment with thionyl chloride to give alternative 3-chloro-8-methyl-8-azabicyclo[3.2.1]octane (XIII) followed by treatment with cyanide as described in J. Am. Chem. Soc., 1958 80, 4677.

As an alternative, compounds of general formula (I) can be prepared from compounds of general formula (VI) by treatment with a suitable base, such as lithium diisopropylamide (LDA), followed by reaction with an aryl or heteroaryl halide (R^1Hal).

Compounds of general formula (VI) can be prepared from 3-cyano-8-azabicyclo[3.2.1]octane (VII) by treatment with a suitable base, such as potassium carbonate, in the presence of an alkyl halide (R^2Hal).

3-Cyano-8-azabicyclo[3.2.1]octane (VII) can be prepared by demethylating 3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (IV) by, for instance, treatment first with a chloroformate ester (such as vinyl chloroformate) to produce a carbamate, followed by acid hydrolysis.

As a further alternative, compounds of general formula (VI) can be prepared by treating compounds of general formula (VIII) with tosylmethyl isocyanide in the presence of a suitable base, such as potassium ethoxide.

Compounds of general formula (VIII) can be prepared by the Robinson tropinone synthesis, see, for instance, J. Chem. Soc., 1917, 111, 762. As an alternative compounds of general formula (VIII) can be prepared from cyclohepta-2,6-dienone (XI) by reaction with an amine (R^3NH_2) as described in, for instance, Tetrahedron, 1973, 155, Bull. Chem. Chem. Soc, Jpn., 1971, 44, 1708 and J. Org. Chem., 1971, 36, 1718.

As yet a further alternative, compounds of general formula (I) can be prepared by treatment of a compound of general formula (IX) with an aryl- or heteroaryl-acetonitrile of general formula (X) in the presence of a suitable base, such as sodium hydride, as described in J. Med. Chem., 1975, 18, 496.

5 The compounds of general formula (VI) (except those where R² represents methyl, benzyl or trichloroethyl are believed not to have been previously described. Accordingly in a further aspect the invention provides compounds of formula (VI) wherein R² has any of the meanings given hereinabove except that R² cannot be methyl, benzyl or trichloroethyl.

10 In a further aspect the invention provides a method of combating insect and like pests at a locus by applying to the locus or the pests an insecticidally-effective amount of an insecticidal composition comprising the compounds of Formula I or an acid addition salt thereof.

15 The compounds of Formula I and acid addition salts thereof may be used to combat and control infestations of insect pests such as Lepidoptera, Diptera, Homoptera and Coleoptera (including Diabrotica i.e. corn rootworms) and also other invertebrate pests, for example, acarine pests. The insect and acarine pests which may be combated and controlled by the use of the invention compounds include those pests associated with agriculture (which term includes the growing of crops for food and fibre products), horticulture and animal husbandry, forestry, the storage of products of vegetable origin, such as fruit, grain and timber, and also
20 those pests associated with the transmission of diseases of man and animals. Examples of insect and acarine pest species which may be controlled by the compounds of Formula I include:

Myzus persicae (aphid), Aphis gossypii (aphid), Aphis fabae (aphid), Aedes aegypti (mosquito), Anopheles spp. (mosquitos), Culex spp. (mosquitos), Dysdercus fasciatus (capsid),
25 Musca domestica (housefly), Pieris brassicae (white butterfly), Plutella xylostella (diamond back moth), Phaedon cochleariae (mustard beetle), Aonidiella spp. (scale insects), Trialeurodes spp. (white flies), Bemisia tabaci (white fly), Blattella germanica (cockroach), Periplaneta americana (cockroach), Blatta orientalis (cockroach) Spodoptera littoralis (cotton leafworm), Heliothis virescens (tobacco budworm) Chortiocetes terminifera (locust),
30 Diabrotica spp. (rootworms), Agrotis spp. (cutworms), Chilo partellus (maize stem borer), Nilaparvata lugens (planthopper), Nephotettix cincticeps (leafhopper), Panonychus ulmi (European red mite), Panonychus citri (citrus red mite), Tetranychus urticae (two-spotted

spider mite), Tetranychus cinnabarinus (carmine spider mite), Phyllocoptura oleivora (citrus rust mite), Polyphagotarsonemus latus (broad mite) and Brevipalpus spp. (mites).

In order to apply the compounds of Formula I to the locus of the nematode, insect or acarid pest, or to a plant susceptible to attack by the nematode, insect or acarid pest, the compound is usually formulated into a composition which includes in addition to the the compounds of Formula I suitable inert diluent or carrier materials, and/or surface active agents. The amount of composition generally applied for the control of nematode pests gives a rate of active ingredient from 0.01 to 10 kg per hectare, preferably from 0.1 to 6 kg per hectare.

The compositions can be applied to the soil, plant or seed, to the locus of the pests, or to the habitat of the pests, in the form of dusting powders, wettable powders, granules (slow or fast release), emulsion or suspension concentrates, liquid solutions, emulsions, seed dressings, fogging/smoke formulations or controlled release compositions, such as microencapsulated granules or suspensions.

Dusting powders are formulated by mixing the active ingredient with one or more finely divided solid carriers and/or diluents, for example natural clays, kaolin, pyrophyllite, bentonite, alumina, montmorillonite, kieselguhr, chalk, diatomaceous earths, calcium phosphates, calcium and magnesium carbonates, sulphur, lime, flours, talc and other organic and inorganic solid carriers.

Granules are formed either by absorbing the active ingredient in a porous granular material for example pumice, attapulgite clays, fuller's earth, kieselguhr, diatomaceous earths, ground corn cobs, and the like, or on to hard core materials such as sands, silicates, mineral carbonates, sulphates, phosphates, or the like. Agents which are commonly used to aid in impregnation, binding or coating the solid carriers include aliphatic and aromatic petroleum solvents, alcohols, polyvinyl acetates, polyvinyl alcohols, ethers, ketones, esters, dextrans, sugars and vegetable oils. with the active ingredient. Other additives may also be included, such as emulsifying agents, wetting agents or dispersing agents.

Microencapsulated formulations (microcapsule suspensions CS) or other controlled release formulations may also be used, particularly for slow release over a period of time, and for seed treatment.

Alternatively the compositions may be in the form of liquid preparations to be used as dips, irrigation additives or sprays, which are generally aqueous dispersions or emulsions of

the active ingredient in the presence of one or more known wetting agents, dispersing agents or emulsifying agents (surface active agents). The compositions which are to be used in the form of aqueous dispersions or emulsions are generally supplied in the form of an emulsifiable concentrate (EC) or a suspension concentrate (SC) containing a high proportion of the active
5 ingredient or ingredients. An EC is a homogeneous liquid composition, usually containing the active ingredient dissolved in a substantially non-volatile organic solvent. An SC is a fine particle size dispersion of solid active ingredient in water. To apply the concentrates they are diluted in water and are usually applied by means of a spray to the area to be treated.

Suitable liquid solvents for ECs include methyl ketone, methyl isobutyl ketone,
10 cyclohexanone, xylenes, toluene, chlorobenzene, paraffins, kerosene, white oil, alcohols, (for example, butanol), methylnaphthalene, trimethylbenzene, trichloroethylene, N-methyl-2-pyrrolidone and tetrahydrofurfuryl alcohol (THFA).

Wetting agents, dispersing agents and emulsifying agents may be of the cationic, anionic or non-ionic type. Suitable agents of the cationic type include, for example,
15 quaternary ammonium compounds, for example cetyltrimethyl ammonium bromide. Suitable agents of the anionic type include, for example, soaps, salts of aliphatic monoesters of sulphuric acid, for example sodium lauryl sulphate, salts of sulphonated aromatic compounds, for example sodium dodecylbenzenesulphonate, sodium, calcium or ammonium lignosulphonate, or butylnaphthalene sulphonate, and a mixture of the sodium salts of
20 diisopropyl- and triisopropylnaphthalene sulphonates. Suitable agents of the non-ionic type include, for example, the condensation products of ethylene oxide with fatty alcohols such as oleyl alcohol or cetyl alcohol, or with alkyl phenols such as octyl phenol, nonyl phenol and octyl cresol. Other non-ionic agents are the partial esters derived from long chain fatty acids and hexitol anhydrides, the condensation products of the said partial esters with ethylene
25 oxide, and the lecithins.

These concentrates are often required to withstand storage for prolonged periods and after such storage, to be capable of dilution with water to form aqueous preparations which remain homogeneous for a sufficient time to enable them to be applied by conventional spray equipment. The concentrates may contain 10-85% by weight of the active ingredient or
30 ingredients. When diluted to form aqueous preparations such preparations may contain varying amounts of the active ingredient depending upon the purpose for which they are to be used.

The compounds of Formula I may also be formulated as powders (dry seed treatment DS or water dispersible powder WS) or liquids (flowable concentrate FS, liquid seed treatment LS, or microcapsule suspension CS) for use in seed treatments.

In use the compositions are applied to the insect pests, to the locus of the pests, to the
5 habitat of the pests, or to growing plants liable to infestation by the pests, by any of the known means of applying pesticidal compositions, for example, by dusting, spraying, or incorporation of granules.

The compound of Formula I may be the sole active ingredient of the composition or they may be admixed with one or more additional active ingredients such as insecticides,
10 synergists, herbicides, fungicides or plant growth regulators where appropriate. Suitable additional active ingredients for inclusion in admixture with a compound of Formula I may be compounds which will broaden the spectrum of activity of the compositions of the invention or increase their persistence in the location of the pest. They may synergise the activity of the compound of Formula I or complement the activity for example by increasing
15 the speed of effect or overcoming repellency. Additionally multi-component mixtures of this type may help to overcome or prevent the development of resistance to individual components. The particular additional active ingredient included will depend upon the intended utility of the mixture and the type of complementary action required. Examples of suitable insecticides include the following:

- 20 a) Pyrethroids such as permethrin, esfenvalerate, deltamethrin, cyhalothrin in particular lambda-cyhalothrin, biphenthrin, fenpropathrin, cyfluthrin, tefluthrin, fish safe pyrethroids for example ethofenprox, natural pyrethrin, tetramethrin, s-bioallethrin, fenfluthrin, prallethrin and 5-benzyl-3-furylmethyl-(E)-(1R,3S)-2,2-dimethyl- 3-(2-oxothiolan-3-ylidenemethyl) cyclopropane carboxylate;
- 25 b) Organophosphates such as profenofos, sulprofos, methyl parathion, azinphos-methyl, demeton-s-methyl, heptenophos, thiometon, fenamiphos, monocrotophos, profenophos, triazophos, methamidophos, dimethoate, phosphamidon, malathion, chloropyrifos, phosalone, terbufos, fensulfothion, fonofos, phorate, phoxim, pyrimiphos-methyl, pyrimiphos-ethyl, fenitrothion or diazinon;
- 30 c) Carbamates (including aryl carbamates) such as pirimicarb, cloethocarb, carbofuran, furathiocarb, ethiofencarb, aldicarb, thiofurox, carbosulfan, bendiocarb, fenobucarb, propoxur or oxamyl;

- 16 -

- d) Benzoyl ureas such as triflumuron, or chlorfluazuron;
 - e) Organic tin compounds such as cyhexatin, fenbutatin oxide, azocyclotin;
 - f) Macrolides such as avermectins or milbemycins, for example such as abamectin, ivermectin, and milbemycin;
 - 5 g) Hormones and pheromones;
 - h) Organochlorine compounds such as benzene hexachloride, DDT, chlordane or dieldrin;
 - i) Amidines, such as chlordimeform or amitraz;
 - j) Fumigant agents;
 - k) Imidacloprid.
- 10 In addition to the major chemical classes of insecticide listed above, other insecticides having particular targets may be employed in the mixture if appropriate for the intended utility of the mixture. For instance selective insecticides for particular crops, for example stemborer specific insecticides for use in rice such as cartap or buprofezin can be employed. Alternatively insecticides specific for particular insect species/stages for example
- 15 ovo-larvicides such as chlofentezine, flubenzimine, hexythiazox and tetradifon, motilicides such as dicofol or propargite, acaricides such as bromopropylate, chlorobenzilate, or growth regulators such as hydramethylnon, cyromazine, methoprene, chlorfluazuron and diflubenzuron may also be included in the compositions.

Examples of suitable synergists for use in the compositions include piperonyl butoxide, 20 sesamax, safroxan and dodecyl imidazole.

Suitable herbicides, fungicides and plant-growth regulators for inclusion in the compositions will depend upon the intended target and the effect required.

An example of a rice selective herbicide which can be included is propanil, an example of a plant growth regulator for use in cotton is "Pix", and examples of fungicides for use in rice 25 include blasticides such as blasticidin-S. The ratio of the compounds of Formula I to the other active ingredient in the composition will depend upon a number of factors including type of target, effect required from the mixture etc. However in general, the additional active ingredient of the composition will be applied at about the rate as it is usually employed, or at a slightly lower rate if synergism occurs.

30 The invention is illustrated by the following examples. Examples 1 to 86 illustrate the preparation of a range of compounds of formula (I).

Examples 87 - 104 illustrate formulations suitable for the application of the compounds of Formula I according to the invention. The following ingredients are referred to by their Registered Trade Marks and have the composition as shown below.

5	<u>Registered Trade Mark Composition</u>	
	Synperonic NP8 }	Nonylphenol-ethylene oxide
	Synperonic NP13 }	condensate
	Synperonic OP10 }	
	Aromasol H	Alkylbenzene solvent
10	Solvesso 200	Inert organic diluent
	Keltrol	Polysaccharide

EXAMPLE 1

This example illustrates the preparation of exo-3-(pyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

15 Potassium t-butoxide (22.4g) was added portionwise to a stirred mixture of tropinone (11.58g) and tosylmethyl isocyanide (21.2g) in dimethoxyethane (240ml) and ethanol (8ml) at 0°C under nitrogen at such a rate to maintain the temperature between 0°C and 10°C. The mixture was then allowed to warm to room temperature and stirred for a further 4 hours. After standing at room
20 temperature for 3 days the mixture was filtered and the solid residue washed with dimethoxyethane. The filtrate was evaporated under reduced pressure and chromatographed [SiO₂; dichloromethane:methanol (90:10)] to give exo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (9.1g).
exo-3-Cyano-8-methyl-8-azabicyclo[3.2.1]octane (10.0g) in tetrahydrofuran (60ml) was added
25 dropwise to a stirred solution of lithium diisopropylamide [made by adding n-BuLi (29ml of a 2.5M solution in hexane) to diisopropylamine (10ml) in tetrahydrofuran (60ml)] at -25°C under nitrogen. The mixture was stirred at -25°C for 20 minutes and then cooled to -78°C. 3-Fluoropyridine (10.0g) in tetrahydrofuran (60ml) was then added dropwise. The mixture was then allowed to warm to room temperature over 6 hours. The mixture was then poured into water and
30 extracted with dichloromethane. The combined extracts were washed with brine, dried (MgSO₄), evaporated under reduced pressure and chromatographed [SiO₂; dichloromethane:methanol (80:20)] to give a yellow oil which crystallised on standing. The solid was washed with hexane

and ether, filtered and air-dried to give exo-3-(pyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (8.2g).

EXAMPLE 2

This example illustrates the preparation of exo-3-(pyrid-3-yl)-endo-3-cyano-8-(2-fluoroethyl)-8-azabicyclo[3.2.1]octane.

Vinyl chloroformate (6.0ml) in tetrahydrofuran (10ml) was added dropwise to exo-3-(pyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (4.0g) in tetrahydrofuran (40ml) at 0°C under nitrogen. The mixture was then heated at 70°C for 4.5 hours. After cooling to room temperature the mixture was filtered and the solid residue washed with ethyl acetate. The combined filtrates were evaporated under reduced pressure and crystallised on standing to give exo-3-(pyrid-3-yl)-endo-3-cyano-8-(vinylloxycarbonyl)-8-azabicyclo[3.2.1]octane (4.1g).

exo-3-(Pyrid-3-yl)-endo-3-cyano-8-(vinylloxycarbonyl)-8-azabicyclo[3.2.1]octane (3.5g) and concentrated hydrochloric acid (3ml) in methanol (25ml) were refluxed for 6 hours and then allowed to stand at room temperature overnight. After the mixture had been refluxed for a further 4 hours it was allowed to cool to room temperature and then evaporated under reduced pressure. The mixture was then partitioned between 2M sodium hydroxide and ethyl acetate and the aqueous layer was separated and extracted with ethyl acetate. The combined organic fractions were washed with brine, dried (MgSO₄) and evaporated under reduced pressure to give exo-3-(pyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (1.7g), which crystallised on standing.

exo-3-(Pyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.2g), 1-bromo-2-fluoroethane (0.21ml), potassium carbonate (0.14g) and tetrahydrofuran (6ml) were heated at 60°C for 6.5 hours and then allowed to stand at room temperature overnight. 1-Bromo-2-fluoroethane (0.2ml) was then added and the mixture heated at 60°C for 6 hours, cooled to room temperature, filtered and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (90:10)] gave exo-3-(pyrid-3-yl)-endo-3-cyano-8-(2-fluoroethyl)-8-azabicyclo[3.2.1]octane (0.123g) m.p. 84.4°C.

EXAMPLE 3

This example illustrates the preparation of exo-3-(3,5-difluorophenyl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

exo-3-Cyano-8-methyl-8-azabicyclo[3.2.1]octane (13.6 g in tetrahydrofuran (80 ml) was added dropwise to a stirred solution of lithium diisopropylamide [made by adding n-BuLi (40ml of a

- 19 -

2.5M solution in hexane) to diisopropylamine (14.0ml) in tetrahydrofuran (80ml)] at -25°C under nitrogen. The mixture was stirred at -25°C for 0.5 hours and then cooled to -78°C. 1,3,5-Trifluorobenzene (12.0g) in tetrahydrofuran (80ml) was added dropwise at such a rate to maintain the temperature below -65°C. The mixture was allowed to warm to room temperature overnight and then poured into water and extracted with dichloromethane. The combined extracts were washed with brine, dried (MgSO₄) and evaporated under reduced pressure to give a yellow solid. This was recrystallised from diethyl ether to give exo-3-(3,5-difluorophenyl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane. The mother liquor from the recrystallisation was chromatographed [SiO₂; dichloromethane:methanol (90:10)] to give further exo-3-(3,5-difluorophenyl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (11.2g in total).

EXAMPLE 4

This example illustrates the preparation of exo-3-(pyrid-3-yl)-endo-3-cyano-8-(prop-1-yl)-8-azabicyclo[3.2.1]octane.

Vinyl chloroformate (2.5ml) in diethyl ether (15ml) was added dropwise to a stirred mixture of exo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (3.0g) in diethyl ether (15ml) at -5°C under nitrogen. The mixture was then stirred at 0°C for 0.5 hours and at reflux for 5 hours. After cooling to room temperature the mixture was filtered and the solid residue washed with diethyl ether. The combined filtrates were evaporated under reduced pressure to give exo-3-cyano-8-(vinylloxycarbonyl)-8-azabicyclo[3.2.1]octane (2.93g).

exo-3-Cyano-8-(vinylloxycarbonyl)-8-azabicyclo[3.2.1]octane (2.9g), concentrated hydrochloric acid (1ml) and methanol (30ml) were refluxed for 4 hours and then allowed to stand at room temperature overnight. Concentrated hydrochloric acid (1ml) was added and the mixture refluxed for 4 hours. After cooling to room temperature the mixture was evaporated under reduced pressure, dissolved in ethyl acetate and washed with 2M sodium hydroxide and brine, dried (MgSO₄) and evaporated under reduced pressure to give exo-3-cyano-8-azabicyclo[3.2.1]octane (1.09g) as a dark yellow solid.

exo-3-Cyano-8-azabicyclo[3.2.1]octane (0.5g), 1-bromopropane (0.34ml) and potassium carbonate (1.27g) were stirred in ethanol (5ml) at room temperature for 5 hours. 1-Bromopropane (0.17ml) was then added and the mixture stirred overnight. 1-Bromopropane (0.17ml) was added and the mixture stirred at room temperature for 6 hours, a further portion of 1-bromopropane (0.17ml) was added and the mixture allowed to stand at room temperature for 3

days and then refluxed for 0.5 hours. The mixture was then cooled to room temperature, filtered and the filtrate evaporated under reduced pressure. Chromatography [SiO_2 ; dichloromethane:methanol (90:10)] gave exo-3-cyano-8-propyl-8-azabicyclo[3.2.1]octane (0.39g). exo-3-Cyano-8-propyl-8-azabicyclo[3.2.1]octane (0.32g) in tetrahydrofuran (92ml) was added dropwise to a stirred solution of lithium diisopropylamide [made by adding n-BuLi (0.8ml of a 2.5M solution in hexane to diisopropylamine (0.2ml) in tetrahydrofuran (2ml)] at -25°C under nitrogen. The mixture was stirred at -25°C for 0.5 hours, cooled to -76°C and 3-fluoropyridine (0.175g) in tetrahydrofuran (2ml) was added dropwise. The mixture was stirred at -76°C for 1 hour and then allowed to warm slowly to room temperature and allowed to stand overnight. The mixture was poured into water, extracted with ethyl acetate (x3) and the combined extracts washed with brine and water, dried (MgSO_4) and evaporated under reduced pressure. Chromatography [SiO_2 ; dichloromethane:methanol (95:5)] gave exo-3-(pyrid-3-yl)-endo-3-cyano-8-(prop-1-yl)-8-azabicyclo[3.2.1]octane (0.35g).

EXAMPLE 5

This example illustrates the preparation of 3-phenyl-3-cyano-8-benzyl-8-azabicyclo[3.2.1]octane.

Sodium hydride (0.75g of a 55% suspension in oil) was carefully added to benzyl cyanide (0.69g) and meso-2,5-bis(chloromethyl)-1-benzylpyrrolidine (1.0g) in N,N-dimethylformamide (30ml) at 0°C under nitrogen. The mixture was stirred at room temperature overnight and then poured into ice-cold water and extracted with dichloromethane. The aqueous layer was allowed to stand at room temperature overnight and then filtered and the solid residue washed with water and air dried. The solid product was chromatographed [SiO_2 ; hexane:ethyl acetate (80:20)] to give a 10:1 (exo-phenyl):(endo-phenyl) mixture of 3-phenyl-3-cyano-8-benzyl-8-azabicyclo[3.2.1]octane (0.21g).

EXAMPLE 6

This example illustrates the preparation of exo-3-(pyrid-3-yl)-endo-3-cyano-8-benzyl-8-azabicyclo[3.2.1]octane.

Three drops of 5M hydrochloric acid was added to a stirred mixture of 2,5-dimethoxytetrahydrofuran (16.5g) and water (70ml). After 10 minutes a mixture of benzylamine (13.6ml) and 5M hydrochloric acid (30ml) were added followed by the immediate addition of a mixture of 1,3-acetonedicarboxylic acid (18.2g) and sodium acetate (10g) in water (100ml). After stirring at room temperature for 3 days, during which carbon dioxide was evolved, the mixture was

- 21 -

basified to pH8 and extracted with ethyl acetate (x3). The combined extracts were dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; hexane:ethyl acetate] to give 8-benzyl-8-azabicyclo[3.2.1]octan-3-one (11.2g).

Potassium t-butoxide (2.5g) was added portionwise to a stirred mixture of 8-benzyl-8-azabicyclo[3.2.1]octan-3-one (2.0g), tosylmethyl isocyanide (2.36g) and ethanol (2ml) in dimethoxyethane (50ml) at 0°C under nitrogen. The mixture was stirred at 0°C for 0.5 hours and then overnight at room temperature. The mixture was then filtered and the solid residue washed with dimethoxyethane. The combined filtrates were evaporated under reduced pressure and chromatographed [SiO₂; hexane:ethyl acetate (80:20)] to give 3-cyano-8-benzyl-8-azabicyclo[3.2.1]octane (0.87g).

3-Cyano-8-benzyl-8-azabicyclo[3.2.1]octane (0.5g) in tetrahydrofuran (2ml) was added dropwise to a stirred solution of lithium diisopropylamide [made by adding n-BuLi (1.5ml of a 1.6M solution in hexane) to diisopropylamine (0.246g) in tetrahydrofuran (2ml)] at -25°C under nitrogen. After 0.5 hours the mixture was cooled to -76°C and 3-fluoropyridine (0.215g) in tetrahydrofuran (2ml) was added. After 2 hours the mixture was allowed to warm room temperature overnight and water then added. The mixture was then extracted with ethyl acetate (x3) and the combined extracts were washed with brine and water, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (95:5)] gave exo-3-(pyrid-3-yl)-endo-3-cyano-8-benzyl-8-azabicyclo[3.2.1]octane (0.245g) which crystallised on standing m.p. 119-120°C.

EXAMPLE 7

This example illustrates the preparation of exo-3-(pyrid-3-yl)-endo-3-cyano-8-(2-methoxyethyl)-8-azabicyclo[3.2.1]octane.

exo-3-(Pyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.30g), 2-bromoethyl methyl ether (0.235g) and potassium carbonate (0.213g) were refluxed in ethanol (3ml) for 30 hours. The mixture was allowed to cool to room temperature, filtered and washed with ethanol. The filtrate was evaporated under reduced pressure and chromatographed [SiO₂; dichloromethane:methanol (95:5)] to give exo-3-(pyrid-3-yl)-endo-3-cyano-8-(2-methoxyethyl)-8-azabicyclo[3.2.1]octane (0.223g).

EXAMPLE 8

This example illustrates the preparation of exo-3-(pyrid-2-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

exo-3-Cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.5g) in tetrahydrofuran (3ml) was added to a solution of lithium diisopropylamide [made by adding n-BuLi (1.6ml of a 2.5M solution in hexane) to diisopropylamine (0.4g) in tetrahydrofuran (3ml)] at -25°C under nitrogen. After 30 minutes a solution of 2-fluoropyridine (0.388g) in tetrahydrofuran (3ml) was added. After 1 hour the mixture was allowed to warm to room temperature and then stand overnight. Water was added and the mixture extracted with ethyl acetate (x3). The combined extracts were washed with water (x2), dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (90:10)] gave exo-3-(pyrid-2-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.467g).

EXAMPLE 9

This example illustrates the preparation of exo-3-(pyrazin-2-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

exo-3-Cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.5g) in tetrahydrofuran (3ml) was added to a solution of lithium diisopropylamide [made by adding n-BuLi (1.6ml of a 2.5M solution in hexane) to diisopropylamine (0.4g) in tetrahydrofuran (3ml)] at -25°C under nitrogen. After 30 minutes the mixture was cooled to -78°C and a solution of chloropyrazine (0.46g) in tetrahydrofuran (5ml) was added. After 1 hour the mixture was allowed to warm to room temperature and stand overnight. Water was added and the mixture extracted with ethyl acetate (x3). The combined extracts were washed with brine and water, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (95:5) to (90:10)] gave exo-3-(pyrazin-2-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.368g) m.p. 76-77°C.

EXAMPLE 10

This example illustrates the preparation of exo-3-(6-chloropyrazin-2-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

exo-3-Cyano-8-methyl-8-azabicyclo[3.2.1]octane (1.0g) in tetrahydrofuran (5ml) was added to a solution of lithium diisopropylamide [made by adding n-BuLi (2.66ml of a 2.5M solution in hexane) to diisopropylamine (0.673g) in tetrahydrofuran (5ml)] at -25°C under nitrogen. After 30 minutes the mixture was cooled to -78°C and a solution of 2,6-dichloropyrazine (1.0g) in tetrahydrofuran (5ml) was added. After 1 hour the mixture was allowed to warm to room temperature and stand over the weekend. Water was added and the mixture extracted with ethyl acetate (x3). The combined extracts were washed with brine and water, dried

(MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (95:5)] gave exo-3-(6-chloropyrazin-2-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (1.10g) m.p. 79.8-80.1°C.

EXAMPLE 11

- 5 This example illustrates the preparation of exo-3-(6-chloropyridazin-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

exo-3-Cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.5g) in tetrahydrofuran (5ml) was added to a solution of lithium diisopropylamide [made by adding n-BuLi (1.4ml of a 2.5M solution in hexane) to diisopropylamine (0.45g) in tetrahydrofuran (2ml)] at -25°C under nitrogen.

- 10 After 30 minutes 1,3-dimethylimidazolidinone (1ml) was added and the mixture cooled to -78°C. A solution of 3,6-dichlorochloropyridazine (0.50g) in tetrahydrofuran (2ml) was added. After 2 hours the mixture was allowed to warm to room temperature and stand overnight. Water was added and the mixture extracted with ethyl acetate (x3). The combined extracts were washed with brine and water, dried (MgSO₄) and evaporated under
15 reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (95:5)] gave exo-3-(6-chloropyridazin-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.082g).

EXAMPLE 12

This example illustrates the preparation of exo-3-(5,6-dichloropyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

- 20 3-Chloro-2-hydroxy-5-nitropyridine (4.8g) was added to phosphorus oxychloride (11ml) and phosphorus pentachloride (4.45g) and the mixture refluxed overnight. The mixture was then cooled to room temperature and evaporated under reduced pressure. Iced water was added to the mixture and a solid product formed. The solid was removed by filtration, washed with water and air-dried to give 2,3-dichloro-5-nitropyridine (3.94g).
- 25 2,3-Dichloro-5-nitropyridine (3.9g) and iron powder (3.0g) were added to isopropyl alcohol (40ml) and water (8ml) and the mixture refluxed for 4 hours. The mixture was then cooled to room temperature and filtered (celite). The filtrate was evaporated under reduced pressure and chromatographed [SiO₂; hexane:ethyl acetate (80:20) to (50:50)] to give 5-amino-2,3-dichloropyridine (1.71g).
- 30 5-Amino-2,3-dichloropyridine (0.80g) in dichloromethane (10ml) was added to boron trifluoride etherate (0.92ml) at -15°C under nitrogen. Dichloromethane (15ml) was added followed by t-butyl nitrite (0.71ml) in dichloromethane (5ml). After 15 minutes the mixture

was allowed to warm to -5°C over 20 minutes. Hexane was added and the resulting solid was filtered, air-dried and washed with ether and stored at approximately -20°C overnight. The solid was then heated until gas evolution had ceased and the product kugelrohr distilled to give 2,3-dichloro-5-fluoropyridine (0.104g).

- 5 exo-3-Cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.10g) in tetrahydrofuran (1ml) was added to a solution of lithium diisopropylamide [made by adding n-BuLi (0.29ml of a 2.5M solution in hexane) to diisopropylamine (0.073g) in tetrahydrofuran (1ml)] at -25°C under nitrogen. After 30 minutes 2,3-dichloro-5-fluoropyridine (0.10g) in tetrahydrofuran (1ml) was added. After 1 hour the mixture was allowed to warm to room temperature and stand overnight.
- 10 Water was added and the resulting mixture extracted with ethyl acetate (x3). The combined extracts were washed with water, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (95:5) to (90:10)] gave an orange gum which was triturated with hexane to give exo-3-(5,6-dichloropyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.019g) as a yellow solid.

15

EXAMPLE 13

This example illustrates the preparation of exo-3-(pyrid-3-yl)-endo-3-cyano-8-(methoxycarbonylmethyl)-8-azabicyclo[3.2.1]octane.

- exo-3-(Pyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.20g), ethyl bromoacetate (0.187g) and potassium carbonate (0.155g) were refluxed in ethanol (3ml) for 4 hours. The
- 20 mixture was then filtered and the filtrate evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (95:5)] gave exo-3-(pyrid-3-yl)-endo-3-cyano-8-(methoxycarbonylmethyl)-8-azabicyclo[3.2.1]octane (0.112g).

EXAMPLE 14

- This example illustrates the preparation of exo-3-(pyrid-3-yl)-endo-3-cyano-8-
- 25 (methylsulphonylmethylsulphonyl)-8-azabicyclo[3.2.1]octane.

exo-3-(Pyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.39g) and triethylamine (15ml) were added to dichloromethane (5ml) and the mixture cooled to -20°C. Methane sulphonyl chloride (0.12ml) was added dropwise and the mixture allowed to warm to room temperature. After 1 hour the mixture was evaporated under reduced pressure and dissolved in ethyl acetate.

The resulting solution was washed with aqueous sodium bicarbonate solution and water (x2), dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (90:10)] gave a gum which formed a solid on trituration with hexane and ether. Chromatography [SiO₂; hexane:ethyl acetate (80:20)] gave exo-3-(pyrid-3-yl)-endo-3-cyano-8-(methylsulphonylmethylsulphonyl)-8-azabicyclo[3.2.1]octane (0.028g) m.p. 163-164°C.

EXAMPLE 15

This example illustrates the preparation of exo-3-(6-chloropyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

10 exo-3-(Pyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.50g) in acetonitrile (3ml) was added dropwise to a stirred solution of di-*t*-butyl carbonate (0.512g) in acetonitrile (5ml) at 0°C. 4-Dimethylaminopyridine (0.02g) was added and after 30 minutes the mixture was warmed to room temperature, stirred for 2 hours and allowed to stand overnight. The mixture was evaporated under reduced pressure and chromatographed [SiO₂; ethyl acetate:dichloromethane (20:80) to (30:70)] to give exo-3-(pyrid-3-yl)-endo-3-cyano-8-(*t*-butyloxycarbonyl)-8-azabicyclo[3.2.1]octane (0.602g).

20 *m*-Chloroperoxybenzoic acid (0.22g) was added to a solution of exo-3-(pyrid-3-yl)-endo-3-cyano-8-(*t*-butyloxycarbonyl)-8-azabicyclo[3.2.1]octane (0.20g) in dichloromethane (2ml) at 0°C under nitrogen. After 1 hour the mixture was warmed to room temperature and allowed to stand overnight. The mixture was evaporated under reduced pressure, dissolved in ethyl acetate, washed with aqueous sodium bicarbonate solution (x2), dried (MgSO₄) and evaporated under reduced pressure to give exo-3-(*N*-oxopyrid-3-yl)-endo-3-cyano-8-(*t*-butyloxycarbonyl)-8-azabicyclo[3.2.1]octane (0.161g).

25 exo-3-(*N*-oxopyrid-3-yl)-endo-3-cyano-8-(*t*-butyloxycarbonyl)-8-azabicyclo[3.2.1]octane (0.161g) was added to phosphorus oxychloride (1ml) and the mixture refluxed for 1 hour. The mixture was then allowed to cool to room temperature, evaporated under reduced pressure, toluene added and evaporated under reduced pressure. Ethyl acetate was added and the mixture washed with aqueous sodium hydroxide solution and water (x2), dried (MgSO₄) and evaporated under reduced pressure to give exo-3-(6-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.058g).

exo-3-(6-Chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.05g) and paraformaldehyde (0.50g) were added to formic acid (2ml) and the mixture heated under reflux. After 2 hours the mixture was allowed to cool to room temperature and stand overnight. The mixture was evaporated under reduced pressure and 2M sodium hydroxide added. The mixture was extracted with ethyl acetate (x3) and the combined extracts were washed with brine and water, dried (MgSO₄), evaporated under reduced pressure and chromatographed [SiO₂; dichloromethane:methanol (95:5)] to give exo-3-(6-chloropyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.023g).

EXAMPLE 16

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(n-hexyl)-8-azabicyclo[3.2.1]octane.

exo-3-Cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.5g) in tetrahydrofuran (3ml) was added to a solution of lithium diisopropylamide [made by adding n-BuLi (1.6ml of a 2.5M solution in hexane) to diisopropylamine (0.4g) in tetrahydrofuran (3ml)] at -25°C under nitrogen. After a further 15 minutes at -25°C 3,5-dichloropyridine (0.588g) in tetrahydrofuran (3ml) was added at -78°C. After 1 hour the mixture was allowed to warm to room temperature and stand overnight. Water was then added and the resulting mixture extracted with ethyl acetate (x3). The combined extracts were washed with brine and water, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (90:10)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.249g).

Vinyl chloroformate (2.6ml) in tetrahydrofuran (5ml) was added to a stirred solution of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (2.6g) in tetrahydrofuran (25ml) at 0°C. The mixture was allowed to warm to room temperature over 1 hour, refluxed for 2 hours and then allowed to cool to room temperature. After 20 hours the mixture was partitioned between water and ethyl acetate and the organic layer was separated, washed with water and dried (MgSO₄). Evaporation under reduced pressure gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(vinylloxycarbonyl)-8-azabicyclo[3.2.1]octane (2.0g).

exo-3-(5-Chloropyrid-3-yl)-endo-3-cyano-8-(vinylloxycarbonyl)-8-azabicyclo[3.2.1]octane (2.6g) was dissolved in methanol (50ml) and concentrated hydrochloric acid (7ml) added. The mixture was refluxed for 3 hours after which the mixture was evaporated under reduced

pressure and basified with aqueous sodium carbonate. The resulting mixture was extracted with ethyl acetate and evaporated under reduced pressure to give a brown solid. This was then washed with hexane to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (1.2g).

5 n-Hexyl bromide (0.1ml) and potassium carbonate (0.1g) were added to exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.15g) in ethanol (2ml) and the mixture refluxed for 44 hours. The mixture was then diluted with ethanol, filtered and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (96:4)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(n-hexyl)-8-azabicyclo[3.2.1]octane
10 (0.123g).

EXAMPLE 17

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-allyl-8-azabicyclo[3.2.1]octane.

Allyl bromide (62μl) and potassium carbonate (0.1g) were added to exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.15g) in ethanol (2ml) and the mixture stirred
15 for 3 hours and then allowed to stand overnight. The mixture was then diluted with ethanol, filtered and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (95:5)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-allyl-8-azabicyclo[3.2.1]octane (0.167g).

20

EXAMPLE 18

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2,2,2-trifluoroethyl)-8-azabicyclo[3.2.1]octane.

A few drops of dilute hydrochloric acid were added to a solution of 2,5-dimethoxytetrahydrofuran (16.5g) in water (70ml). After stirring at room temperature for 30
25 minutes 2,2,2-trifluoroethylamine hydrochloride (16.9g), 1,3-acetonedicarboxylic acid (18.3g) and sodium acetate (10.0g) were added and the mixture stirred at room temperature for 2 days. The mixture was diluted to 500ml with water, saturated with potassium carbonate and extracted with ethyl acetate (x2). The combined organic extracts were washed with aqueous potassium carbonate, dried (MgSO₄) and evaporated under reduced pressure. Distillation
30 (90°C; 0.1mmHg) gave 8-(2,2,2-trifluoroethyl)-8-azabicyclo[3.2.1]octan-3-one (8.7g).

Potassium t-butoxide (5.4g) was added slowly with cooling to a stirred solution of 8-(2,2,2-trifluoroethyl)-8-azabicyclo[3.2.1]octan-3-one (4.0g) and tosylmethyl isocyanide (4.9g) in

dimethoxyethane (80ml) and ethanol (5ml) under nitrogen at such a rate so as to keep the temperature below 10°C. The mixture was stirred for 18 hours while allowing it to warm to room temperature, evaporated under reduced pressure and added to aqueous potassium carbonate solution. The mixture was extracted with ethyl acetate (x2) and the combined
5 extracts were dried (MgSO₄) and evaporated under reduced pressure to give an oil. The mixture was extracted with refluxing hexane and the extracts allowed to cool and evaporated under reduced pressure to give exo-3-cyano-8-(2,2,2-trifluoroethyl)-8-azabicyclo[3.2.1]octane (2.5g) m.p. 90-92°C.

exo-3-Cyano-8-(2,2,2-trifluoroethyl)-8-azabicyclo[3.2.1]octane (1.09g) in tetrahydrofuran
10 (10ml) was added to a stirred solution of lithium diisopropylamide [made by adding n-BuLi (2.4ml of a 2.5M solution in hexane) to diisopropylamine (0.61g) in tetrahydrofuran (10ml)] at -25°C under nitrogen. After 2 hours at -25°C the mixture was cooled to -76°C and 3,5-dichloropyridine (0.74g) in tetrahydrofuran (10ml) added. The mixture was allowed to warm to room temperature, stirred for 18 hours and evaporated under reduced pressure. The
15 mixture was dissolved in ether, washed with water (x2), dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; diethyl ether:hexane (20:80) to (50:50)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2,2,2-trifluoroethyl)-8-azabicyclo[3.2.1]octane (0.45g) m.p. 109.5-111.5°C.

EXAMPLE 19

20 This example illustrates the preparation of exo-3-(5-bromopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.
exo-3-Cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.5g) in tetrahydrofuran (3ml) was added to a solution of lithium diisopropylamide [made by adding n-BuLi (1.6ml of a 2.5M solution in hexane) to diisopropylamine (0.4g) in tetrahydrofuran (3ml)] at -25°C under nitrogen.
25 After 30 minutes the mixture was cooled to -76°C and a solution of 3,5-dibromopyridine (0.94g) in tetrahydrofuran (3ml) added. After 1 hour the mixture was allowed to warm to room temperature and left to stand overnight. Water was added and the resulting mixture extracted with ethyl acetate (x3). The combined extracts were washed with brine (x2) and water, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂;
30 dichloromethane:methanol (90:10)] gave exo-3-(5-bromopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.327g) m.p. 144-145°C.

EXAMPLE 20

This example illustrates the preparation of exo-3-(5-cyanopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

exo-3-(5-Bromopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.30g) and
5 copper(I) cyanide (0.345g) were heated at 200°C in N-methylpyrrolidinone (10ml) under
nitrogen. After 36 hours the reaction was allowed to cool to room temperature and water
was added followed by aqueous ammonium hydroxide solution (density=0.88). The mixture
was extracted with ethyl acetate (x3) and the combined extracts were washed with brine and
water, dried (MgSO₄) and evaporated under reduced pressure. The resulting oil was
10 dissolved in ether and washed with brine (x7), dried (MgSO₄) and evaporated under reduced
pressure. Chromatography [SiO₂; dichloromethane:methanol (95:5)] gave a yellow solid.
This was recrystallised three times (from dichloromethane/hexane, ethyl acetate/hexane and
dichloromethane/hexane) to give exo-3-(5-cyanopyrid-3-yl)-endo-3-cyano-8-methyl-8-
azabicyclo[3.2.1]octane (0.49g) m.p. 183.5-184°C.

15

EXAMPLE 21

This example illustrates the preparation of exo-3-(5-ethoxypyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

exo-3-(5-Chloropyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.30g) and
20 sodium ethoxide (0.625g) were heated at 80°C in N,N-dimethylformamide (10ml) under
nitrogen. After 5 hours the mixture was allowed to cool to room temperature and water
added. The mixture was extracted with ethyl acetate (x3) and the combined extracts were
washed with brine (x2) and water, dried (MgSO₄) and evaporated under reduced pressure.
Chromatography [SiO₂; dichloromethane:methanol (90:10)] gave an oil. A small amount of
25 hexane was added and the mixture was allowed to stand at approximately 0°C overnight after
which a solid product had formed. The mixture was filtered and the solid washed with a small
amount of hexane to give exo-3-(5-ethoxypyrid-3-yl)-endo-3-cyano-8-methyl-8-
azabicyclo[3.2.1]octane (0.105g) m.p. 56-57°C.

EXAMPLE 22

30 This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-isopropyl-8-azabicyclo[3.2.1]octane.

- 2M Hydrochloric acid (8 drops) was added to a stirred solution of 2,5-dimethoxytetrahydrofuran (16.5g) in water (70ml). After 15 minutes a mixture of diisopropylamine (7.38g) and 2M hydrochloric acid (40ml) was added to the reaction followed by acetonedicarboxylic acid (18.25g) and sodium acetate (10.0g) in water (100ml).
- 5 After 3 days 1,3-acetonedicarboxylic acid (6.0g) and sodium acetate (3.0g) were added. After a further 6 days the mixture was basified to pH8 and extracted with ethyl acetate. The extracts were dried (MgSO₄) and evaporated under reduced pressure. The aqueous fraction was then extracted with chloroform and the extracts dried (MgSO₄) and evaporated under reduced pressure. Distillation of the combined extracts (95-115°C; 18mmHg) gave 8-isopropyl-8-azabicyclo[3.2.1]octan-3-one (3.37g).
- 10 Potassium t-butoxide (5.0g) was added slowly with cooling to a stirred solution of 8-isopropyl-8-azabicyclo[3.2.1]octan-3-one (3.16g) and tosylmethyl isocyanide (4.80g) in dimethoxyethane (50ml) and ethanol (2.2ml) under nitrogen at such a rate so as to keep the temperature below 10°C. After 1 day tosylmethyl isocyanide (1.0g), potassium t-butoxide (1.0g) and ethanol (1ml) were added. After a further day the mixture was filtered and the filtrate evaporated under reduced pressure and chromatographed [SiO₂; dichloromethane:methanol (95:5)] to give exo-3-cyano-8-isopropyl-8-azabicyclo[3.2.1]octane (0.90g).
- 15
- 20 Lithium bis(trimethylsilyl)amide (2.5ml of a 1M solution in tetrahydrofuran) in tetrahydrofuran (5ml) was added to a stirred solution of exo-3-cyano-8-isopropyl-8-azabicyclo[3.2.1]octane (0.38g) and 3,5-dichloropyridine (0.34g) in tetrahydrofuran (5ml) at 10°C over 30 minutes. The mixture was then stirred at room temperature for 2 hours and allowed to stand at room temperature overnight. 3,5-Dichloropyridine (0.15g) was added
- 25 followed by lithium bis(trimethylsilyl)amide (1.0ml of a 1M solution in tetrahydrofuran) over 30 minutes. After 2 hours lithium bis(trimethylsilyl)amide (1.0ml of a 1M solution in tetrahydrofuran) was added dropwise and after a further 1 hour additional lithium bis(trimethylsilyl)amide (1.0ml of a 1M solution in tetrahydrofuran) was added and the mixture warmed to 50°C. After 5 minutes the reaction was cooled to room temperature and
- 30 aqueous sodium carbonate solution added. The mixture was extracted with ethyl acetate (x2) and the combined extracts washed with brine, dried (MgSO₄) and evaporated under reduced pressure to give a brown oil. The oil was extracted with boiling hexane and the combined

extracts evaporated under reduced pressure to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-isopropyl-8-azabicyclo[3.2.1]octane (0.60g).

EXAMPLE 23

This example illustrates the preparation of exo-3-(2,6-dichloropyrimid-4-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

exo-3-Cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.5g) in tetrahydrofuran (3ml) was added to a solution of lithium diisopropylamide [made by adding n-BuLi (1.6ml of a 2.5M solution in hexane) to diisopropylamine (0.4g) in tetrahydrofuran (3ml)] at -25°C under nitrogen. After 30 minutes the mixture was cooled to -78°C and a solution of 2,4,6-trichloropyrimidine (0.728g) in tetrahydrofuran (5ml) was added. After 1 hour the mixture was allowed to warm to room temperature, stirred for 2 hours and allowed to stand overnight. Water was added and the mixture extracted with ethyl acetate (x3). The combined extracts were washed with brine and water, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (95:5) to (90:10)] gave exo-3-(2,6-dichloropyrimid-4-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.087g) m.p. 95-97°C.

EXAMPLE 24

This example illustrates the preparation of exo-3-(2-chloropyrid-4-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

exo-3-Cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.5g) in tetrahydrofuran (3ml) was added to a solution of lithium diisopropylamide [made by adding n-BuLi (1.6ml of a 2.5M solution in hexane) to diisopropylamine (0.4g) in tetrahydrofuran (3ml)] at -25°C under nitrogen. After 30 minutes the mixture was cooled to -78°C and a solution of 2,4,6-trichloropyridine (0.724g) in tetrahydrofuran (5ml) was added. After 1 hour the mixture was allowed to warm to room temperature and allowed to stand overnight. Water was added and the mixture extracted with ethyl acetate (x3). The combined extracts were dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (95:5)] gave a solid product which was recrystallised (ethyl acetate/hexane) to give exo-3-(2,6-dichloropyrid-4-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.389g) m.p. 165-166°C.

exo-3-(2,6-Dichloropyrid-4-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.50g) and hydrazine hydrate (0.106ml) were refluxed in isopropyl alcohol (5ml) for 5 hours and then left to stand overnight. Hydrazine hydrate (0.106ml) was added and the mixture refluxed

for 8 hours. More hydrazine hydrate (0.106ml) was added and the mixture refluxed for a further 8 hours. After cooling to room temperature the mixture was evaporated under reduced pressure and the residue extracted with dichloromethane. The extracts washed with water (x2), dried (MgSO₄), evaporated under reduced pressure and triturated with hexane
5 and ether to give exo-3-(2-chloro-6-hydrazinopyrid-4-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.205g) m.p. 215-216°C.

Copper(II) sulphate octahydrate (0.36g) was added to a solution of exo-3-(2-chloro-6-hydrazinopyrid-4-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.170g) in water (3ml) and the mixture refluxed for 7 hours. After cooling to room temperature ammonium
10 hydroxide solution (density=0.88) was added and the mixture extracted with ethyl acetate (x3). The combined extracts were washed with brine and water, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (90:10)] gave exo-3-(2-chloropyrid-4-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.052g) m.p. 104-105°C.

15

EXAMPLE 25

This example illustrates the preparation of exo-3-(pyrid-4-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

exo-3-(2,6-Dichloropyrid-4-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.35g),
crushed potassium hydroxide (0.133g) and palladium on charcoal (0.20g) were stirred in
20 methanol (10ml) under hydrogen for 3 hours and then allowed to stand for 3 days. The mixture was filtered (celite), evaporated under reduced pressure and dissolved in ethyl acetate. The resulting solution was washed with aqueous sodium hydroxide and water, dried (MgSO₄) and evaporated under reduced pressure to give a solid product which was washed with hexane and ether to give exo-3-(pyrid-4-yl)-endo-3-cyano-8-methyl-8-
25 azabicyclo[3.2.1]octane (0.072g) m.p. 74.5-76°C.

EXAMPLE 26

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(3,3-difluoroprop-2-en-1-yl)-8-azabicyclo[3.2.1]octane.

exo-3-(5-Chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.248g), 1-bromo-1,1-
30 difluoroprop-2-ene (0.314g) and potassium carbonate (0.345g) were stirred in ethanol (2ml) for 2 hours and then allowed to stand for 4 days. The mixture was then evaporated under reduced pressure and water added. The mixture was then extracted with dichloromethane

(x3) and the combined extracts were washed with brine, dried (MgSO₄) and evaporated under reduced pressure. Filtration [SiO₂; dichloromethane:methanol (98:2)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(3,3-difluoroprop-2-en-1-yl)-8-azabicyclo[3.2.1]octane (0.287g).

5

EXAMPLE 27

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(3-oxo-4,4,4-trifluorobut-1-en-1-yl)-8-azabicyclo[3.2.1]octane.

exo-3-(5-Chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.30g), 4-ethoxy-1,1,1-trifluorobut-3-en-2-one (0.204g) and potassium carbonate (0.20g) were heated under reflux
10 in ethanol. After 4 hours the mixture was allowed to cool to room temperature and water added. The mixture was extracted with ethyl acetate (x3) and the combined extracts were washed with brine (x2) and water, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (95:5)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(3-oxo-4,4,4-trifluorobut-1-en-1-yl)-8-azabicyclo[3.2.1]octane (0.162g) m.p.
15 144-145°C.

EXAMPLE 28

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-acetyl-8-azabicyclo[3.2.1]octane.

N,N-Diisopropylethylamine (0.43ml) and acetyl chloride (0.18ml) were added to exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.50g) in dichloromethane (10ml)
20 at room temperature. After 10 minutes the mixture was evaporated under reduced pressure and ethyl acetate (50ml) added. The resulting mixture was washed with potassium carbonate solution, dried (MgSO₄) and evaporated under reduced pressure. The resulting product was triturated with hot hexane and evaporated under reduced pressure to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-acetyl-8-azabicyclo[3.2.1]octane (0.43g) m.p. 162-165°C.
25

EXAMPLE 29

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane hydroperchlorate.

Perchloric acid (1.19ml) was added dropwise to a stirred suspension of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (5.0g) in diethyl ether (100ml) at room
30 temperature. After 5 hours the mixture was filtered and the precipitate washed with diethyl

ether to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane hydroperchlorate (5.36g).

EXAMPLE 30

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(tert-butyl)-8-azabicyclo[3.2.1]octane.

Acetone (0.42ml) was added to a stirred solution of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane hydroperchlorate (1.0g) in ethanol (2ml) at room temperature under nitrogen. After 30 minutes the mixture was heated to 50°C. After 1 hour the mixture was evaporated under reduced pressure. Acetone was then added and the mixture heated under reflux for 3 hours and then evaporated under reduced pressure. Diethyl ether (10ml) was added followed by methylmagnesium bromide (4.3ml of a 3.0M solution in diethyl ether). The mixture was then heated under reflux for 6 hours and then allowed to stand at room temperature overnight. Saturated ammonium chloride solution was then added and the mixture extracted with dichloromethane (x3). The combined extracts were washed with brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (95:5)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(tert-butyl)-8-azabicyclo[3.2.1]octane (0.218g) m.p. 127-129°C.

EXAMPLE 31

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenyl-3-oxo-prop-2-yl)-8-azabicyclo[3.2.1]octane.

2-Phenylpropanal (1.08g) was added to a mixture of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (2.0g) and p-toluenesulphonic acid (0.15g) in toluene (30ml) and the mixture heated under Dean and Stark reflux for 3 hours. After standing at room temperature overnight the mixture was evaporated under reduced pressure to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenylprop-1-en-1-yl)-8-azabicyclo[3.2.1]octane which was used without further purification.

Sodium N-chloro-p-toluenesulphonamide (2.3g) was added to the exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenylprop-1-en-1-yl)-8-azabicyclo[3.2.1]octane from the above reaction in dichloromethane (30ml) and the mixture stirred at room temperature for 5 hours. After standing at room temperature over the weekend the mixture was stirred for 8 hours and then allowed to stand overnight. The mixture was then filtered (celite) and the residue washed with dichloromethane. The combined filtrates were washed with sodium hypochlorite (x2)

- 35 -

and brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (98:2)] followed by chromatography [SiO₂; dichloromethane:methanol (99:1)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenyl-3-oxo-prop-2-yl)-8-azabicyclo[3.2.1]octane (0.43g) m.p. 124-126°C.

5

EXAMPLE 32

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenylbut-3-en-2-yl)-8-azabicyclo[3.2.1]octane.

Sodium methoxide (0.085g) was added in two portions to a stirred solution of methyltriphenylphosphonium bromide (0.56g) in dimethyl sulphoxide (30ml) at room temperature under nitrogen. The mixture was warmed to 70°C and after 2 hours exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenyl-3-oxo-prop-2-yl)-8-azabicyclo[3.2.1]octane (0.30g) in a small volume of dimethyl sulphoxide was added dropwise. After 3 hours the mixture was allowed to cool to room temperature and stand overnight. The mixture was poured into ice/water and the resulting mixture extracted with ethyl acetate (x3). The combined extracts were washed with brine (x2), dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; ethyl acetate:hexane (90:10)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenylbut-3-en-2-yl)-8-azabicyclo[3.2.1]octane (0.24g).

15

EXAMPLE 33

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenyl-3-hydroxyprop-2-yl)-8-azabicyclo[3.2.1]octane.

20

Sodium borohydride (0.094g) was added to a stirred solution of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenyl-3-oxo-prop-2-yl)-8-azabicyclo[3.2.1]octane (0.90g) in ethanol (15ml) under nitrogen. After 2 hours the mixture was poured into brine and the resulting mixture extracted with ethyl acetate (x2). The combined extracts were dried (MgSO₄), evaporated under reduced pressure and chromatographed [SiO₂; ethyl acetate:hexane (50:50)] to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenyl-3-hydroxyprop-2-yl)-8-azabicyclo[3.2.1]octane (0.777g) m.p. 124-126°C.

25

EXAMPLE 34

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-fluoro-2-phenylprop-1-yl)-8-azabicyclo[3.2.1]octane.

30

Diethylaminosulphur trifluoride (0.4ml) and exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenyl-3-hydroxyprop-2-yl)-8-azabicyclo[3.2.1]octane (0.10g) were stirred in

- 36 -

dichloromethane (0.2ml) at room temperature for 4 hours. The mixture was allowed to stand at room temperature over the weekend and water added. The mixture was extracted with ethyl acetate and the aqueous layer basified with saturated sodium bicarbonate solution. The aqueous layer was then extracted with ethyl acetate (x3) and the combined organic extracts
5 were washed with sodium bicarbonate solution and brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; ethyl acetate:hexane (17:83)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-fluoro-2-phenylprop-1-yl)-8-azabicyclo[3.2.1]octane (0.075g).

EXAMPLE 35

10 This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenyl-3-acetoxyp-2-yl)-8-azabicyclo[3.2.1]octane.
Triethylamine (0.06ml) and acetyl chloride (0.029ml) were added to a stirred solution of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenyl-3-hydroxyprop-2-yl)-8-azabicyclo[3.2.1]octane (0.15g) in dichloromethane (5ml) at room temperature under
15 nitrogen. After 1.5 hours dichloromethane was added and the mixture washed with water (x2) and brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; ethyl acetate:hexane (20:80)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenyl-3-acetoxyp-2-yl)-8-azabicyclo[3.2.1]octane (0.14g) m.p. 130-131°C.

EXAMPLE 36

20 This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-formyl-8-azabicyclo[3.2.1]octane.
exo-3-(5-Chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (3.0g) and formic acid (1.14ml) were heated at reflux for 4 hours. The mixture was then heated at 110°C overnight and formic acid (1.0ml) added. After 8 hours the mixture was allowed to cool to room
25 temperature and stand overnight. Ethyl acetate was added and the mixture washed with 2M sodium hydroxide solution (x2), water and brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; ethyl acetate:methanol (95:5)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-formyl-8-azabicyclo[3.2.1]octane (1.675g) m.p. 141-142°C.

30

EXAMPLE 37

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(diisopropylcarbamy)-8-azabicyclo[3.2.1]octane.

Triethylamine (0.27ml) followed by diisopropylcarbonyl chloride (0.317g) was added to a stirred solution of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.40g) in dichloromethane (5ml) at room temperature. After 2 hours the mixture was allowed to stand at room temperature for 4 days. Dichloromethane was then added and the mixture washed with water (x2) and brine, dried (MgSO₄) and evaporated under reduced pressure. Filtration [SiO₂; ethyl acetate] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(diisopropylcarbonyl)-8-azabicyclo[3.2.1]octane (0.12g) m.p. 118-121°C.

EXAMPLE 38

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(tert-butylcarbonyl)-8-azabicyclo[3.2.1]octane.

Triethylamine (0.27ml) followed by tert-butylisocyanate (0.22ml) was added to a stirred solution of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.40g) in dichloromethane (4ml) at room temperature. After 3 hours the mixture was allowed to stand at room temperature overnight and dichloromethane then added. The mixture was then washed with water (x2) and brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; diethyl ether] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(tert-butylcarbonyl)-8-azabicyclo[3.2.1]octane (0.40g) m.p. 62-65°C.

EXAMPLE 39

This example illustrates the preparation of (R)-exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(1-phenylethyl)-8-azabicyclo[3.2.1]octane.

Three drops of 5M hydrochloric acid were added to a mixture of 2,5-dimethoxytetrahydrofuran (16.5g) and water (70ml). A cooled mixture of (R)- α -methylbenzylamine (15.125g) and 5M hydrochloric acid (30ml) was then added followed by 1,3-acetonedicarboxylic acid (18.26g) sodium acetate (10g) and water (100ml). After 5 days the mixture was basified with aqueous sodium carbonate solution and extracted with ethyl acetate. The extracts were washed with brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; ethyl acetate:hexane (10:90) to (20:80)] gave (R)-8-(1-phenylethyl)-8-azabicyclo[3.2.1]octan-3-one.

Potassium *t*-butoxide (13.4g) was added portionwise to a stirred mixture of (R)-8-(1-phenylethyl)-8-azabicyclo[3.2.1]octan-3-one (11.45g) and tosylmethyl isocyanide (12.7g) in dimethoxyethane (200ml) and ethanol (6ml) at -5°C at such a rate to maintain the temperature below -2°C. After stirring overnight the mixture was filtered (celite) and the filtrate evaporated

under reduced pressure. The residue was then dissolved in ethyl acetate, washed with water, dried (MgSO_4) and evaporated under reduced pressure. Chromatography [SiO_2 ; ethyl acetate:hexane (50:50)] gave (R)-exo-3-cyano-8-(1-phenylethyl)-8-azabicyclo[3.2.1]octane (3.5g) m.p. 138-139.5°C.

- 5 Lithium bis(trimethylsilyl)amide (4.8ml of a 1.0M solution in tetrahydrofuran) was added to a stirred solution of (R)-exo-3-cyano-8-(1-phenylethyl)-8-azabicyclo[3.2.1]octane (1.0g) and 3,5-dichloropyridine (0.674g) in tetrahydrofuran (20ml) at 0°C under nitrogen. The mixture was allowed to warm to room temperature and stand for 24 hours. Water (20ml) was added and the mixture stirred for 30 minutes and then allowed to stand for 2 days. The mixture was
10 extracted with ethyl acetate and the extracts washed with brine, dried (MgSO_4) and evaporated under reduced pressure. Chromatography [SiO_2 ; ethyl acetate:hexane (50:50)], preparative thin layer chromatography [SiO_2 ; ethyl acetate:hexane (25:75)] and recrystallisation from hexane gave (R)-exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(1-phenylethyl)-8-azabicyclo[3.2.1]octane (0.46g) m.p. 113-115°C.

15

EXAMPLE 40

This example illustrates the preparation of exo-3-(5-aminopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

- Ammonia solution (35%) was added to exo-3-(5-bromopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.106g) and copper(II) sulphate hydrate (0.001g) and the tube
20 sealed. The mixture was heated at 100°C for 20 hours and then 150°C for 24 hours. The mixture was then cooled and evaporated under reduced pressure. The residue was then dissolved in methanol, charcoal added and the mixture filtered and evaporated under reduced pressure. Water and dichloromethane were added followed by ammonia solution and the resulting mixture was extracted with dichloromethane. The combined extracts were washed
25 with brine, dried (MgSO_4) and evaporated under reduced pressure to give exo-3-(5-aminopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.045g) m.p. 188-190°C.

EXAMPLE 41

- This example illustrates the preparation of exo-3-(5-acetylamidopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.
30

Acetic anhydride (1.0ml) was added to exo-3-(5-aminopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.10g). After 3 days dilute sodium bicarbonate solution and ethyl

acetate were added followed by sodium bicarbonate and potassium carbonate to basify the mixture. The mixture was extracted with ethyl acetate and the extracts dried (MgSO₄) and evaporated under reduced pressure to give exo-3-(5-acetylamidopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.107g).

5

EXAMPLE 42

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(a-cyanobenzyl)-8-azabicyclo[3.2.1]octane.

exo-3-(5-Chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (2.0g) and benzaldehyde (0.89ml) were added to 1M hydrochloric acid (20ml) and the mixture stirred for 20 minutes.

10 Sodium cyanide (0.549g) in water (6ml) was then added. After 18 hours ethanol (20ml) was added to give one phase. After 6 days the reaction mixture was partitioned between ethyl acetate and water and the organic layer was washed with water and brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol:triethylamine (99.4:0.5:0.1)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(a-cyanobenzyl)-8-azabicyclo[3.2.1]octane (0.104g) m.p. 141-142°C.

15

EXAMPLE 43

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(a-carbamylbenzyl)-8-azabicyclo[3.2.1]octane.

Concentrated sulphuric acid (10ml) was added to exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(a-cyanobenzyl)-8-azabicyclo[3.2.1]octane (0.51g) and the mixture stirred for 1 hour. Ice (100g) was added and the mixture basified with sodium bicarbonate solution. A precipitate formed which was collected by filtration, dissolved in ethyl acetate, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (99:1) to (98:2)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(a-carbamylbenzyl)-8-azabicyclo[3.2.1]octane (0.181g) m.p. 193-195°C.

20

25

EXAMPLE 44

This example illustrates the preparation of exo-3-(5-iodopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

Nickel(II) bromide (1.55ml of a 0.16M solution in N,N-dimethylformamide) was added to a stirred solution of tri(n-butyl)phosphine (0.124ml) in N,N-dimethylformamide (5ml) under nitrogen. Potassium iodide (3.96g) was then added followed by exo-3-(5-bromopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (1.522g) and the mixture heated under

30

reflux for 48 hours. The mixture was then cooled to room temperature and partitioned between water and ethyl acetate. The organic layer was dried (MgSO₄), evaporated under reduced pressure and chromatographed [SiO₂; dichloromethane:methanol (95:5)] to give exo-3-(5-iodopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.399g) m.p. 144-145°C.

EXAMPLE 45

This example illustrates the preparation of exo-3-(5-trifluoromethylpyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

exo-3-(5-iodopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.50g) followed by copper(I) iodide were added to a stirred solution of sodium trifluoroacetate (2.6g) in N-methylpyrrolidinone (5ml) and the mixture heated to 180°C. After 3 hours the mixture was cooled to room temperature, water added and extracted with dichloromethane. The organic layer was filtered and the filtrate dried (MgSO₄) and evaporated under reduced pressure. Diethyl ether was added and the mixture extracted repeatedly with water. The aqueous fraction was evaporated under reduced pressure, basified with potassium carbonate and extracted with diethyl ether. The extracts were washed with 1M hydrochloric acid and the aqueous fraction basified with potassium carbonate and extracted with diethyl ether. The extracts were dried (MgSO₄) and evaporated under reduced pressure. Preparative thin layer chromatography [Al₂O₃; diethyl ether] gave exo-3-(5-trifluoromethylpyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.027g) m.p. 118.2-118.5°C.

EXAMPLE 46

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(mercaptothiocarbonyl)-8-azabicyclo[3.2.1]octane.

Carbon disulphide (0.12ml) was added to a stirred solution of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.50g) in ethanol (5ml) at room temperature under nitrogen. After 4 hours the mixture was allowed to stand overnight. The precipitate was collected by filtration and washed with hexane to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(mercaptothiocarbonyl)-8-azabicyclo[3.2.1]octane (0.509g) m.p. 224°C (decomposed).

EXAMPLE 47

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(fluorocarbonyl)-8-azabicyclo[3.2.1]octane.

Iodomethane (0.08ml) was added to a stirred mixture of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(mercaptothiocarbonyl)-8-azabicyclo[3.2.1]octane (0.40g) in dimethyl sulphoxide (3ml) at room temperature. After 3 hours water was added and the mixture extracted with dichloromethane (x3). The combined extracts were washed with water (x2) and brine, dried
5 (MgSO₄) and evaporated under reduced pressure. Dichloromethane was added and the mixture washed with brine (x2), dried (MgSO₄) and evaporated under reduced pressure to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(methylmercaptothiocarbonyl)-8-azabicyclo[3.2.1]octane (0.187g).

Tetra(n-butyl)ammonium dihydrogentrifluoride (0.48g) was added to a solution of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(methylmercaptothiocarbonyl)-8-azabicyclo[3.2.1]octane
10 (0.180g) in dichloromethane at 0°C under nitrogen. N-Bromosuccinimide (0.38g) was then added. After 10 minutes the mixture was warmed to room temperature. After 2 hours the mixture was cooled to 0°C and allowed to stand over the weekend. The mixture was then stirred at room temperature for 8 hours and allowed to stand overnight. The mixture was
15 diluted with dichloromethane and sodium bicarbonate and sodium bisulphite solutions added. The mixture was extracted with dichloromethane and the extracts dried (MgSO₄) and evaporated under reduced pressure. Ethyl acetate was added and the mixture washed with water (x2) and brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; diethyl ether:hexane (80:20)] gave exo-3-(5-chloropyrid-3-yl)-endo-
20 3-cyano-8-(fluorocarbonyl)-8-azabicyclo[3.2.1]octane (0.10g) m.p. 165-167°C.

EXAMPLE 48

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2,2-difluoroethyl)-8-azabicyclo[3.2.1]octane.

2,2-Difluoroethyl bromide (1.08g), potassium carbonate (1.38g), potassium iodide (0.30g)
25 and exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (1.238g) were stirred at 50°C in ethanol (10ml). After 8 hours the mixture was allowed to stand at room temperature for 3 days. 2,2-Difluoroethyl bromide (1.08g), was then added and the mixture heated under reflux for 48 hours. 2,2-Difluoroethyl bromide (1.08g) and potassium carbonate (1.38g) were then added and the mixture heated under reflux for 24 hours. 2,2-Difluoroethyl
30 bromide (1.08g) was then added and the mixture heated under reflux for 24 hours. The mixture was then cooled to room temperature and water added. The mixture was extracted with dichloromethane (x2) and the combined extracts washed with brine, dried (MgSO₄) and

evaporated under reduced pressure. Chromatography [SiO_2 ; dichloromethane:methanol (96:4)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2,2-difluoroethyl)-8-azabicyclo[3.2.1]octane (0.278g) m.p. 101-104°C.

EXAMPLE 49

- 5 This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenylethyl)-8-azabicyclo[3.2.1]octane.
- 2-Phenylethyl bromide (0.222g), potassium carbonate (0.345g) and exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.248g) were heated under reflux in ethanol (2ml) for 9 hours. 2-Phenylethyl bromide (0.1g) was added and the mixture refluxed for 5
- 10 hours. The mixture was then cooled to room temperature and water added. The mixture was extracted with dichloromethane (x2) and the combined extracts washed with brine, dried (MgSO_4) and evaporated under reduced pressure. Chromatography [SiO_2 ; dichloromethane:methanol (98:2)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenylethyl)-8-azabicyclo[3.2.1]octane (0.08g).

15

EXAMPLE 50

- This example illustrates the preparation of exo-3-(5-hydroxypyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.
- Pyridinium hydrochloride (1.0g) and exo-3-(5-methoxypyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.20g) were heated together at 150°C for 5 hours. The mixture was
- 20 then cooled to room temperature, water added and the mixture basified with sodium bicarbonate solution and extracted with ethyl acetate (x3). The aqueous fraction was neutralised with dilute hydrochloric acid and extracted with ethyl acetate (x3). The combined organic extracts were washed with water, dried (MgSO_4) and evaporated under reduced pressure. Chromatography [SiO_2 ; dichloromethane:methanol (90:10) to (80:20)] gave a gum
- 25 which crystallised on addition of diethyl ether to give exo-3-(5-hydroxypyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.049g) m.p. 171-172°C.

EXAMPLE 51

- This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-benzyl-8-azabicyclo[3.2.1]octane.
- 30 exo-3-Cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.5g) in tetrahydrofuran (3ml) was added to a solution of lithium diisopropylamide [made by adding n-BuLi (1.6ml of a 2.5M solution in hexane) to diisopropylamine (0.4g) in tetrahydrofuran (3ml)] at -25°C under nitrogen.

After 15 minutes at the mixture was cooled to -78°C and 3,5-dichloropyridine (0.588g) in tetrahydrofuran (3ml) was added. After 1 hour the mixture was allowed to warm to room temperature and stand overnight. Water was then added and the resulting mixture extracted with ethyl acetate (x3). The combined extracts were washed with brine and water, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (90:10)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.249g).

Vinyl chloroformate (2.6ml) in tetrahydrofuran (5ml) was added to a stirred solution of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (2.6g) in tetrahydrofuran (25ml) at 0°C. The mixture was allowed to warm to room temperature over 1 hour, refluxed for 2 hours and then allowed to cool to room temperature. After 20 hours the mixture was partitioned between water and ethyl acetate and the organic layer was separated, washed with water and dried (MgSO₄). Evaporation under reduced pressure gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(vinylloxycarbonyl)-8-azabicyclo[3.2.1]octane (2.0g).

exo-3-(5-Chloropyrid-3-yl)-endo-3-cyano-8-(vinylloxycarbonyl)-8-azabicyclo[3.2.1]octane (2.6g) was dissolved in methanol (50ml) and concentrated hydrochloric acid (7ml) added. The mixture was refluxed for 3 hours after which the mixture was evaporated under reduced pressure and basified with aqueous sodium carbonate. The resulting mixture was extracted with ethyl acetate and evaporated under reduced pressure to give a brown solid. This was then washed with hexane to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (1.2g).

exo-3-(5-Chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.104g) in ethanol (5ml) was added to benzyl bromide (0.079g) and potassium carbonate (0.12g) and the mixture refluxed for 18 hours. After cooling to room temperature the mixture was evaporated under reduced pressure and partitioned between water and ethyl acetate. The organic layer was separated and evaporated under reduced pressure. Preparative thin layer chromatography [SiO₂; dichloromethane:methanol (97:3)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-benzyl-8-azabicyclo[3.2.1]octane (0.077g).

30

EXAMPLE 52

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(pentafluorophenylmethyl)-8-azabicyclo[3.2.1]octane.

exo-3-(5-Chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.248g), 2,3,4,5,6-pentafluorobenzyl bromide (0.313g), potassium carbonate (0.345g) and ethanol (2ml) were stirred under reflux for 3 hours. The mixture was then evaporated under reduced pressure and water added. The mixture was then extracted with dichloromethane (x3) and the
5 combined extracts were washed with brine, dried (MgSO₄) and evaporated under reduced pressure to give an oil which crystallised on standing. The crystals were washed with a small volume of ether to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(pentafluorophenylmethyl)-8-azabicyclo[3.2.1]octane (0.258g) m.p. 143-144°C.

EXAMPLE 53

10 This example illustrates the preparation of potassium exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(4-carboxylatobenzyl)-8-azabicyclo[3.2.1]octane.

exo-3-(5-Chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.248g) 4-bromomethylbenzoic acid (0.258g), potassium carbonate (0.345g) and ethanol (2ml) were stirred under reflux for 2.5 hours. The mixture was then diluted with ethanol, filtered and the
15 filtrate evaporated under reduced pressure to give potassium exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(4-carboxylatobenzyl)-8-azabicyclo[3.2.1]octane (0.292g).

EXAMPLE 54

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(3-chloro-4-fluorobenzyl)-8-azabicyclo[3.2.1]octane.

20 exo-3-(5-Chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.495g), 3-chloro-4-fluorobenzaldehyde (0.317g) and formic acid (96%, 0.230g) were heated under reflux for 5 hours. The mixture was then cooled to room temperature, basified with dilute sodium hydroxide and extracted with dichloromethane (x2). The combined extracts were washed with brine, dried (MgSO₄), and evaporated under reduced pressure. Chromatography [SiO₂;
25 dichloromethane:methanol (100:0) to (95:5)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(3-chloro-4-fluorobenzyl)-8-azabicyclo[3.2.1]octane (0.290g) m.p. 95-97°C.

EXAMPLE 55

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(pyrid-2-ylmethyl)-8-azabicyclo[3.2.1]octane.

30 2-Picolyl chloride hydrochloride (0.361g), potassium carbonate (0.828g) and exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.495g) were heated under reflux in ethanol (4ml) for 2 hours. The mixture was then cooled to room temperature and water

added. The mixture was extracted with dichloromethane (x2) and the combined extracts washed with brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (100:0) to (95:5)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(pyrid-2-ylmethyl)-8-azabicyclo[3.2.1]octane (0.447g) m.p. 123-125°C.

5

EXAMPLE 56

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-((2-methylthiazol-4-yl)methyl)-8-azabicyclo[3.2.1]octane.

4-Chloromethyl-2-methylthiazole hydrochloride (0.202g), potassium carbonate (0.483g) and exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.247g) were heated under reflux in ethanol (2ml) for 1.5 hours. 4-Chloromethyl-2-methylthiazole hydrochloride (0.40g) was then added and the mixture refluxed for 30 minutes. The mixture was then cooled to room temperature and water added. The mixture was extracted with dichloromethane (x2) and the combined extracts washed with brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (99:1) to (95:5)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-((2-methylthiazol-4-yl)methyl)-8-azabicyclo[3.2.1]octane (0.269g) m.p. 81-83°C.

15

EXAMPLE 57

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-((3,5-dimethylisoxazol-4-yl)methyl)-8-azabicyclo[3.2.1]octane.

4-Chloromethyl-3,5-dimethylisoxazole (0.160g), potassium carbonate (0.345g), potassium iodide (0.02g) and exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.247g) were heated under reflux in ethanol (2ml) for 3 hours. The mixture was then cooled to room temperature and water added. The mixture was extracted with dichloromethane (x2) and the combined extracts washed with brine, dried (MgSO₄) and evaporated under reduced pressure. Filtration [SiO₂; dichloromethane:methanol (98:2)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-((3,5-dimethylisoxazol-4-yl)methyl)-8-azabicyclo[3.2.1]octane (0.258g) m.p. 95-99°C.

25

EXAMPLE 58

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(5-chlorothiophen-2-yl)-8-azabicyclo[3.2.1]octane.

30

2-Chloro-5-chloromethylthiophene (0.367g), potassium carbonate (0.690g) and exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.495g) were heated under reflux

in ethanol (2ml) for 1.5 hours. Potassium iodide (0.02g) was then added and the mixture refluxed for 1.5 hours. The mixture was then cooled to room temperature and water added. The mixture was extracted with dichloromethane (x2) and the combined extracts washed with brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (100:0) to (96:4)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(5-chlorothiophen-2-yl)-8-azabicyclo[3.2.1]octane (0.37g) m.p. 119-121°C.

EXAMPLE 59

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-((5-chloro-1,2,3-thiadiazol-4-yl)methyl)-8-azabicyclo[3.2.1]octane.

5-Chloro-4-chloromethyl-1,2,3-thiadiazole (0.187g), potassium carbonate (0.345g) and exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.248g) were heated under reflux in ethanol (2ml) for 4 hours. The mixture was then cooled to room temperature and water added. The mixture was extracted with dichloromethane (x2) and the combined extracts washed with brine, dried (MgSO₄) and evaporated under reduced pressure.

Chromatography [SiO₂; dichloromethane:methanol (100:0) to (98:2)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-((5-chloro-1,2,3-thiadiazol-4-yl)methyl)-8-azabicyclo[3.2.1]octane (0.148g).

EXAMPLE 60

This example illustrates the preparation of exo-3-(5-fluoropyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

exo-3-(5-Aminopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.40g) in dichloromethane (150ml) was added to boron trifluoride etherate (1.5ml) at -10 to -15°C. After a few minutes *t*-butyl nitrite (2ml) was added and the mixture allowed to warm to room temperature and stand overnight. The solid precipitate was collected and heated to cause decomposition. The residue was dissolved in 2M hydrochloric acid, washed with ethyl acetate, basified and extracted with ethyl acetate. The extracts were dried (MgSO₄), evaporated under reduced pressure and chromatographed [SiO₂; dichloromethane:methanol (90:10)] to give exo-3-(5-fluoropyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.089g).

EXAMPLE 61

This example illustrates the preparation of exo-3-(5-(pyrrol-1-yl)pyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

2,5-Dimethoxytetrahydrofuran (0.53ml) was added to a mixture of exo-3-(5-aminopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane and acetic acid (13ml). After 5 minutes the mixture was heated at reflux for 1 hour and then allowed to cool to room temperature and stand overnight. Ethyl acetate was added and the mixture extracted with 2M hydrochloric acid and water. The combined extracts were basified with potassium carbonate and extracted with ethyl acetate. The extracts were dried (MgSO₄) and evaporated under reduced pressure to give exo-3-(5-(pyrrol-1-yl)pyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.105g).

EXAMPLE 62

10 This example illustrates the preparation of exo-3-(5-(1-ethoxyvinyl)pyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

(1-Ethoxyvinyl)tri-n-butyltin ((0.82ml) was added to a stirred mixture of exo-3-(5-iodopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.817g) and N,N-dimethylformamide (30ml) at room temperature under nitrogen. Bis(triphenylphosphine)palladium(II) chloride (0.65g) was then added and the mixture heated at 130°C for 3 hours. The mixture was then allowed to cool to room temperature, water added and the mixture extracted with dichloromethane (x3). The combined extracts were washed with brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (91:9)] followed by chromatography [SiO₂; dichloromethane:methanol (98:2) to (92:8)] gave exo-3-(5-(1-ethoxyvinyl)pyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.22g).

EXAMPLE 63

This example illustrates the preparation of exo-3-(5-acetylpyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

2M Hydrochloric acid (1ml) was added to a stirred mixture of exo-3-(5-(1-ethoxyvinyl)pyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.18g) in acetone (2ml). After 3 hours the mixture was allowed to stand overnight. The mixture was poured onto saturated sodium bicarbonate solution and the resulting mixture extracted with dichloromethane (x3). The combined extracts were dried (MgSO₄), evaporated under reduced pressure and chromatographed [SiO₂; dichloromethane:methanol (93:7)] to give exo-3-(5-acetylpyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.13g).

EXAMPLE 64

This example illustrates the preparation of exo-3-(5-ethynylpyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

exo-3-(5-Iodopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.50g) in
5 tetrahydrofuran (1ml) was added dropwise to a stirred mixture of trimethylsilylacetylene
(0.22ml), diethylamine (1.13ml), copper(I) iodide (0.01g) and
tetrakis(triphenylphosphine)palladium(0) (0.02g) at room temperature under nitrogen. After
3 hours the mixture was allowed to stand at room temperature for 24 hours and then
evaporated under reduced pressure. Dichloromethane (10ml) was added followed by
10 tetrabutylammonium fluoride (1.7ml of a 1M solution in tetrahydrofuran) and the mixture
stirred at room temperature for 1.5 hours. Water was added and the mixture extracted with
dichloromethane (x2). The combined extracts were washed with brine, dried (MgSO₄),
evaporated under reduced pressure and chromatographed [SiO₂; dichloromethane:methanol
(95:5) to (90:10)] to give a crude product. Chromatography [SiO₂;
15 dichloromethane:methanol (92:8)] followed by dissolving the product in dichloromethane,
washing with water (x2) and brine, drying (MgSO₄) and evaporating under reduced pressure
gave exo-3-(5-ethynylpyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.40g).

EXAMPLE 65

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-
20 (1-hydroxy-1-cyano-2-phenylprop-2-yl)-8-azabicyclo[3.2.1]octane.
Sodium cyanoborohydride (0.033g) was added to a stirred solution of isopropylamine
(0.045ml) and exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenyl-3-oxo-prop-2-yl)-8-
azabicyclo[3.2.1]octane (0.20g) in methanol (2ml). Methanolic hydrogen chloride was then
added to give pH5. After 2 hours the mixture was allowed to stand at room temperature for
25 2 days and isopropylamine (0.045ml) and sodium cyanoborohydride (0.033g) were added.
After stirring at room temperature for 6 hours saturated sodium bicarbonate was added and
the mixture extracted with dichloromethane (x3). The combined extracts were washed with
brine, dried (MgSO₄), evaporated under reduced pressure and chromatographed [SiO₂;
dichloromethane:methanol (100:0) to (90:10)] to give exo-3-(5-chloropyrid-3-yl)-endo-3-
30 cyano-8-(1-hydroxy-1-cyano-2-phenylprop-2-yl)-8-azabicyclo[3.2.1]octane (0.75g) m.p. 172-
175°C.

EXAMPLE 66

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-(isopropylamino)-2-phenylprop-1-yl)-8-azabicyclo[3.2.1]octane.

Triethylamine (0.1ml) and methanesulphonyl chloride (0.053ml) were added to a stirred
5 solution of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-phenyl-3-hydroxyprop-2-yl)-8-azabicyclo[3.2.1]octane (0.25g) in dichloromethane (5ml) at room temperature. After 2 hours isopropylamine (0.65ml) was added. After 2 hours the mixture was allowed to stand at room temperature overnight and dichloromethane then added. The mixture was washed with water (x2) and brine, dried (MgSO₄), evaporated under reduced pressure and
10 chromatographed [SiO₂; dichloromethane:methanol (98:2) to (95:5)] to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-(isopropylamino)-2-phenylprop-1-yl)-8-azabicyclo[3.2.1]octane (0.24g).

EXAMPLE 67

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-
15 cyanomethyl-8-azabicyclo[3.2.1]octane.

Bromoacetonitrile (0.85ml) was added to a mixture of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (2.0g) and potassium carbonate (2.23g) in ethanol (10ml) and the mixture heated at reflux for 3 hours. The mixture was then cooled to room temperature, filtered (celite) and washed through with dichloromethane. The filtrates were
20 evaporated under reduced pressure, chromatographed [SiO₂; dichloromethane:methanol (99:1)] and recrystallised (ethyl acetate/hexane) to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-cyanomethyl-8-azabicyclo[3.2.1]octane (1.36g). m.p. 149-151°C.

EXAMPLE 68

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(1-
25 (ethoxycarbonyl)ethyl)-8-azabicyclo[3.2.1]octane.

Ethyl 2-bromopropionate (0.29ml) was added to a mixture of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.5g) and potassium carbonate (0.42g) in tetrahydrofuran (8ml) and the mixture heated at reflux for 24 hours. The mixture was then cooled to room temperature, filtered (celite) and washed through with dichloromethane. The
30 filtrates were evaporated under reduced pressure and chromatographed [SiO₂; dichloromethane:methanol (100:0 to (98:2)] to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(1-(ethoxycarbonyl)ethyl)-8-azabicyclo[3.2.1]octane (0.49g).

EXAMPLE 69

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(6-fluoropyrid-2-yl)-8-azabicyclo[3.2.1]octane.

2,6-Difluoropyridine (0.37ml), potassium carbonate (1.12g) and exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (1.0g) were heated at 140°C in N-methylpyrrolidinone (10ml) for a total of 8 hours. The mixture was then cooled to room temperature, poured into water and the resulting mixture extracted with ethyl acetate (x3). The combined extracts were washed with brine, dried (MgSO₄), evaporated under reduced pressure and chromatographed [SiO₂; hexane:ethyl acetate (90:10)] to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(6-fluoropyrid-2-yl)-8-azabicyclo[3.2.1]octane (0.796g) m.p. 131.5-132.5°C.

EXAMPLE 70

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(5-chlorothiazol-2-yl)-8-azabicyclo[3.2.1]octane.

2-Bromo-5-chlorothiazole (2.4g), potassium carbonate (1.67g) and exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (1.0g) were heated at 140°C in N-methylpyrrolidinone (10ml) for a total of 10 hours. The mixture was then cooled to room temperature, poured into water and the resulting mixture extracted with ethyl acetate (x3). The combined extracts were washed with brine, dried (MgSO₄), evaporated under reduced pressure and chromatographed [SiO₂; hexane:ethyl acetate (90:10)]. Recrystallisation (hexane) gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(5-chlorothiazol-2-yl)-8-azabicyclo[3.2.1]octane (0.17g) m.p. 111-112°C.

EXAMPLE 71

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-pentafluorophenyl-8-azabicyclo[3.2.1]octane.

Hexafluorobenzene (0.93ml), potassium carbonate (1.12g) and exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (1.0g) were heated at 150°C in N-methylpyrrolidinone (10ml) for 5 hours. Hexafluorobenzene (0.93ml) and potassium carbonate (1.12g) were added and the mixture heated at 160°C for 7 hours. The mixture was then cooled to room temperature, poured into water and the resulting mixture extracted with ethyl acetate (x3). The combined extracts were washed with brine, dried (MgSO₄),

evaporated under reduced pressure and chromatographed [SiO_2 ; hexane:ethyl acetate (90:10)] to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-pentafluorophenyl-8-azabicyclo[3.2.1]octane (0.55g).

EXAMPLE 72

5 This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-cyano-8-azabicyclo[3.2.1]octane.

Tosyl cyanide (0.88ml) was added dropwise to a stirred mixture of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (1.0g) and *N,N*-diisopropylethylamine (0.85ml) in tetrahydrofuran (5ml) at room temperature. After 6 hours the mixture was allowed to stand at room temperature overnight, poured into water and extracted with ethyl acetate (x3). The combined extracts were washed with water and brine, dried (MgSO_4), evaporated under reduced pressure and chromatographed [SiO_2 ; hexane:ethyl acetate (90:10)]. Recrystallisation (hexane/ethyl acetate) gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-cyano-8-azabicyclo[3.2.1]octane (0.20g) m.p. 168-170°C.

15

EXAMPLE 73

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-methoxy-8-azabicyclo[3.2.1]octane.

N,N-Diisopropylethylamine (14.5ml) was added dropwise to a stirred suspension of *O*-methylhydroxylamine hydrochloride (2.32g) in isopropyl alcohol (25ml). After 30 minutes cyclohepta-2,6-dienone (3.0g) in isopropyl alcohol (5ml) was added dropwise. After 24 hours *N,N*-diisopropylethylamine (4.9ml) was added. After 6 hours the mixture was allowed to stand at room temperature overnight. The mixture was evaporated under reduced pressure, diethyl ether added and the resulting mixture extracted with 2*M* hydrochloric acid (x3). The combined aqueous fractions were washed with diethyl ether (x3), neutralised with sodium hydroxide and extracted with diethyl ether (x3). The combined extracts were washed with brine, dried (MgSO_4) and evaporated under reduced pressure. Kugelrohr distillation gave 8-methoxy-8-azabicyclo[3.2.1]octan-3-one (0.86g).

Tosylmethyl isocyanide (2.52g) was added to a stirred suspension of potassium *t*-butoxide (2.17g) in 1,2-dimethoxyethane (10ml) at such a rate to keep the temperature below 10°C. After 45 minutes 8-methoxy-8-azabicyclo[3.2.1]octan-3-one (1.0g) in 1,2-dimethoxyethane (10ml) was added dropwise. After 30 minutes the mixture was allowed to warm to room temperature. After 4 hours the mixture was allowed to stand at room temperature overnight

30

and water was then added. The resulting mixture was extracted with ethyl acetate (x3) and the combined extracts washed with brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; hexane:ethyl acetate (90:10)] gave exo-3-cyano-8-methoxy-8-azabicyclo[3.2.1]octane (0.40g).

- 5 Lithium bis(trimethylsilyl)amide (2.42ml of a 1M solution in tetrahydrofuran) was added dropwise to a stirred solution of exo-3-cyano-8-methoxy-8-azabicyclo[3.2.1]octane (0.40g) and 3,5-dichloropyridine (0.358g) in tetrahydrofuran (5ml) at 0°C. After 1 hour the mixture was allowed to warm to room temperature. After 5 hours water was added and the mixture extracted with ethyl acetate (x3). The combined extracts were washed with brine, dried
10 (MgSO₄) and evaporated under reduced pressure. Preparative thin layer chromatography [SiO₂; ethyl acetate] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-methoxy-8-azabicyclo[3.2.1]octane (0.192g) m.p. 107.5-108.5°C.

EXAMPLE 74

- This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-
15 (ethoxycarbonyl)ethyl)-8-azabicyclo[3.2.1]octane.

- Sodium hydride (0.095g of an 80% dispersion in oil) was added to exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.75g) and ethyl acrylate (2.0g) in tetrahydrofuran. The mixture was refluxed for 8 hours then allowed to cool to room temperature, water added and the mixture extracted with ethyl acetate (x2). The combined
20 extracts were washed with brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; chloroform:methanol (95:5)] followed by chromatography [SiO₂; ethyl acetate:dichloromethane (80:20)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-(ethoxycarbonyl)ethyl)-8-azabicyclo[3.2.1]octane.

EXAMPLE 75

- 25 This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-carboxyethyl)-8-azabicyclo[3.2.1]octane.

- 3M Sodium hydroxide (4ml) was added to a stirred solution of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-(ethoxycarbonyl)ethyl)-8-azabicyclo[3.2.1]octane (0.41g) in ethanol (8ml) at room temperature. After 24 hours the mixture was basified to pH9 and evaporated under
30 reduced pressure. The product was azeotroped with methanol/toluene and chromatographed [SiO₂; dichloromethane:methanol (75:25)] to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-carboxyethyl)-8-azabicyclo[3.2.1]octane (0.21g) m.p. 180-181°C.

EXAMPLE 76

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(Q,Q-diethylphosphonomethyl)-8-azabicyclo[3.2.1]octane.

exo-3-(5-Chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.20g), Q,Q-diethylphosphonomethyl triflate (0.245g) and potassium carbonate (0.15g) were heated under reflux in tetrahydrofuran (8ml). After 3 hours the mixture was cooled to room temperature, filtered and evaporated under reduced pressure. Chromatography [SiO_2 ; dichloromethane:methanol (96:4)] followed by high pressure liquid chromatography [SiO_2 ; dichloromethane:methanol (96:4)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(Q,Q-diethylphosphonomethyl)-8-azabicyclo[3.2.1]octane m.p. 69-70°C.

EXAMPLE 77

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-phosphonomethyl-8-azabicyclo[3.2.1]octane.

Trimethylsilyl bromide (1.5ml) was added dropwise to a stirred solution of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(Q,Q-diethylphosphonomethyl)-8-azabicyclo[3.2.1]octane (0.56g) in dichloromethane (30ml) at 0°C. After 30 minutes the mixture was allowed to warm to room temperature. After 7 hours trimethylsilyl bromide (0.8ml) was added, after 23 hours more trimethylsilyl bromide (0.5ml) was added and after 18 hours further trimethylsilyl bromide (0.5ml) was added. After 24 hours the mixture was evaporated under reduced pressure, water added and the mixture filtered. After 10 minutes the filtrate was azeotroped with methanol/toluene to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-phosphonomethyl-8-azabicyclo[3.2.1]octane (0.49g) m.p. 242-245°C.

EXAMPLE 78

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-cyanoethyl)-8-azabicyclo[3.2.1]octane.

3-Bromopropionitrile (0.174ml) was added to exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.40g) and potassium carbonate (0.45g) in ethanol (10ml) and the mixture heated under reflux for 16 hours. 3-Bromopropionitrile (0.13ml) was added and the mixture refluxed for 3 hours and allowed to cool to room temperature. Water was added and the mixture extracted with dichloromethane (x3). The combined extracts were washed with brine, dried (MgSO_4) and evaporated under reduced pressure. Chromatography [SiO_2 ;

dichloromethane:methanol (100:0) to (98:2) gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-cyanoethyl)-8-azabicyclo[3.2.1]octane (0.343g).

EXAMPLE 79

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(1,1-dimethyl-2,2,2-trifluoroethyl)-8-azabicyclo[3.2.1]octane.

4Å Molecular sieves (1.0g) were added to a suspension of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane hydroperchlorate (3.42g) in acetone (30ml) and the mixture heated under reflux for 5 hours. The mixture was then allowed to cool to room temperature and filtered to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-isopropylene-8-azabicyclo[3.2.1]octanium perchlorate (0.279g).

Trimethyl(trifluoromethyl)silane (5.2ml of a 0.5M solution in tetrahydrofuran) was added to a suspension of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-isopropylene-8-azabicyclo[3.2.1]octanium perchlorate (0.50g) in tetrahydrofuran (5ml). Cesium fluoride (0.39g) was added and the mixture placed in an ultrasound bath for 2.5 hours. The mixture was then added to water and the resulting mixture extracted with dichloromethane (x3). The combined extracts were washed with brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (97:3)] followed by preparative thin layer chromatography [SiO₂; dichloromethane:methanol (98:2)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(1,1-dimethyl-2,2,2-trifluoroethyl)-8-azabicyclo[3.2.1]octane (0.02g).

EXAMPLE 80

This example illustrates the preparation of exo-3-(5-(2,2,2-trifluoroethoxy)pyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane.

Sodium (0.46g) was added portionwise to a solution of 2,2,2-trifluoroethanol (2.3ml) in *N*-methylpyrrolidinone (20ml) under nitrogen. Tetraphenylphosphonium bromide (0.05g) and exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (2.6g) were added and the mixture heated at 110°C for 18 hours and 140°C for 5 hours. Sodium (0.6g) was added to a solution of 2,2,2-trifluoroethanol (3ml) in *N*-methylpyrrolidinone (5ml) and after 30 minutes the resulting mixture was added to the reaction mixture. After 6 hours at 140°C the mixture was cooled to room temperature and added to water. The mixture was extracted with diethyl ether (x2) and the combined extracts were washed with water, dried (MgSO₄) and evaporated under reduced pressure. The resulting mixture was filtered [SiO₂;

dichloromethane:methanol (95:5)] and chromatographed [SiO₂; dichloromethane:methanol (95:5)] to give exo-3-(5-(2,2,2-trifluoroethoxy)pyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.065g, 80% pure).

EXAMPLE 81

5 This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(1-cyanoethyl)-8-azabicyclo[3.2.1]octane.

exo-3-(5-Chloropyrid-3-yl)-endo-3-cyano-8-azabicyclo[3.2.1]octane (0.50g), 2-bromopropionitrile (2ml) and potassium carbonate (0.50g) were refluxed in ethanol (5ml). After 24 hours the mixture was cooled to room temperature, water added and the mixture
10 extracted with dichloromethane (x3). The combined extracts were washed with brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (100:0) to (98:2)] followed by chromatography [SiO₂; ethyl acetate:hexane (80:20)] and preparative thin layer chromatography [Al₂O₃; ethyl acetate:hexane (40:60)] gave exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(1-cyanoethyl)-8-
15 azabicyclo[3.2.1]octane (0.044g, 80% pure).

EXAMPLE 82

This example illustrates the preparation of exo-3-(5-phenylpyrid-3-yl)-endo-3-cyano-8-(vinylloxycarbonyl)-8-azabicyclo[3.2.1]octane.

Vinyl chloroformate (2.17ml) was added a solution of exo-3-(5-iodopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (3.0g) in tetrahydrofuran (20ml) at 0°C. The
20 mixture was then heated under reflux for 5 hours and then stand at room temperature overnight. Water was then added and the mixture extracted with dichloromethane (x3). The combined extracts were washed with water and brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; dichloromethane:methanol (98:2)] gave exo-3-(5-iodopyrid-3-yl)-endo-3-cyano-8-(vinylloxycarbonyl)-8-azabicyclo[3.2.1]octane (3.64g).
25

Tetrakis(triphenylphosphine)palladium(0) (0.042g) was added to a stirred solution of exo-3-(5-iodopyrid-3-yl)-endo-3-cyano-8-(vinylloxycarbonyl)-8-azabicyclo[3.2.1]octane (0.50g) in toluene (2ml). To this mixture was added 2M sodium carbonate solution (1.22ml) and phenylboronic acid (0.16g) in ethanol (0.5ml) and the mixture heated under reflux. After 2
30 hours the mixture was cooled to room temperature, water added and the mixture extracted with ethyl acetate (x3). The combined extracts were washed with brine, dried (MgSO₄) and evaporated under reduced pressure. Chromatography [SiO₂; ethyl acetate:hexane (45:55)]

gave exo-3-(5-phenylpyrid-3-yl)-endo-3-cyano-8-(vinylloxycarbonyl)-8-
azabicyclo[3.2.1]octane (0.33g).

EXAMPLE 83

This example illustrates the preparation of exo-3-(5-phenylpyrid-3-yl)-endo-3-cyano-8-
5 azabicyclo[3.2.1]octane.

Concentrated hydrochloric acid (0.5ml) was added to a solution of exo-3-(5-phenylpyrid-3-yl)-endo-3-cyano-8-(vinylloxycarbonyl)-8-
azabicyclo[3.2.1]octane (0.30g) in methanol (10ml) and the mixture heated under reflux. After 5 hours the mixture was allowed to stand at room
temperature for 4 days and then heated under reflux for 5 hours. The mixture was then
10 allowed to cool to room temperature, basified with saturated sodium bicarbonate solution and
extracted with dichloromethane (x3). The combined extracts were washed with brine and
extracted with 2M hydrochloric acid (x2). The acidic extracts were basified and re-extracted
with dichloromethane (x3). The combined organic extracts were washed with brine, dried
(MgSO₄) and evaporated under reduced pressure to give exo-3-(5-phenylpyrid-3-yl)-endo-3-
15 cyano-8-azabicyclo[3.2.1]octane (0.20g).

EXAMPLE 84

This example illustrates the preparation of exo-3-(5-methylpyrid-3-yl)-endo-3-cyano-8-
(vinylloxycarbonyl)-8-azabicyclo[3.2.1]octane.

Methylolithium (8.7ml of a 1.4M solution in diethyl ether) was added dropwise to a stirred
20 suspension of copper(I) iodide (1.16g) in diethyl ether (10ml) at 0°C under nitrogen. After 45
minutes exo-3-(5-iodopyrid-3-yl)-endo-3-cyano-8-(vinylloxycarbonyl)-8-
azabicyclo[3.2.1]octane (0.50g) in diethyl ether (5ml) was added. After 5 days at room
temperature water was added and the mixture extracted with ethyl acetate (x3). The
combined extracts were washed with brine, dried (MgSO₄) and evaporated under reduced
25 pressure. Chromatography [SiO₂; ethyl acetate:hexane (60:40)] gave exo-3-(5-methylpyrid-
3-yl)-endo-3-cyano-8-(vinylloxycarbonyl)-8-azabicyclo[3.2.1]octane (0.085g).

EXAMPLE 85

This example illustrates the preparation of exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-
cyanoprop-2-yl)-8-azabicyclo[3.2.1]octane.

30 Sodium cyanide (0.069g) was added to a stirred solution of exo-3-(5-chloropyrid-3-yl)-endo-
3-cyano-8-isopropylene-8-azabicyclo[3.2.1]octanium perchlorate (0.50g) in acetonitrile (5ml)
at room temperature under nitrogen. After 4 hours the mixture was allowed to stand at room

temperature over the weekend, water added and the mixture extracted with dichloromethane (x3). The combined extracts were washed with brine, dried (MgSO₄) and evaporated under reduced pressure to give exo-3-(5-chloropyrid-3-yl)-endo-3-cyano-8-(2-cyanoprop-2-yl)-8-azabicyclo[3.2.1]octane (0.38g, 90% pure).

5

EXAMPLE 86

This example illustrates the preparation of exo-3-(5-(ethoxycarbonyl)pyrid-3-yl)-endo-3-cyano-8-(vinylloxycarbonyl)-8-azabicyclo[3.2.1]octane.

Potassium carbonate (0.205g) and bis(triphenylphosphine)palladium(II) chloride (0.026g) were added to a stirred solution of exo-3-(5-iodopyrid-3-yl)-endo-3-cyano-8-methyl-8-azabicyclo[3.2.1]octane (0.50g) in ethanol (10ml) under nitrogen. The reaction vessel was then flushed with carbon monoxide, triethylamine (3 drops) added and the mixture heated under reflux. After 3 hours the mixture was cooled to room temperature, water and brine added and the mixture extracted with ethyl acetate (x3). The combined extracts were dried (MgSO₄), evaporated under reduced pressure and chromatographed [SiO₂; ethyl acetate:hexane (35:65)] to give exo-3-(5-(ethoxycarbonyl)pyrid-3-yl)-endo-3-cyano-8-(vinylloxycarbonyl)-8-azabicyclo[3.2.1]octane (0.29g).

Confirmation of the structural identity of the compounds prepared in Examples 20 to 106 was obtained by proton magnetic resonance spectroscopy. The results are set out in the following table.

20

EXAMPLE

¹H NMR (270MHz), in CDCl₃ unless otherwise stated

- | | |
|---|---|
| 1 | 8.81 (1H, d), 8.56 (1H, dd), 7.85 (1H, dt), 7.30 (1H, dd), 3.44 (2H, m), 2.45-2.15 (8H, m) and 2.33 (3H, s). |
| 2 | 8.80 (1H, d), 8.55 (1H, dd), 7.85 (1H, m), 7.30 (1H, m), 4.55 (2H, dt), 3.4 (2H, m), 2.70 (2H, dt) and 2.4-2.1 (8H, m). |
| 3 | 7.09 (2H, m), 6.75 (1H, m), 3.31 (2H, m), 2.4-2.1 (8H, m) and 2.35 (3H, s). |
| 4 | 8.80 (1H, d), 8.54 (1H, dd), 7.83 (1H, dt), 7.29 (1H, dd), 3.40 (2H, m), 2.4-2.0 (10H, m), 1.48 (2H, hex) and 0.91 (3H, s). |
| 5 | 7.53 (2H, m), 7.4-7.2 (8H, m), 3.59 (2H, s), 3.36 (2H, m) and 2.5-2.15 (8H, m). |
| 6 | 8.82 (1H, d), 8.58 (1H, dd), 7.86 (1H, dt), 7.4-7.2 (6H, m), 3.59 (2H, s), 3.39 (2H, m) and 2.45-2.15 (8H, m). |
| 7 | 8.80 (1H, d), 8.56 (1H, dd), 7.84 (1H, dt), 7.31 (1H, dd), 3.50 (2H, t), 3.48 (2H, m), 3.36 (3H, s), 2.61 (2H, t) and 2.4-2.05 (8H, m). |
| 8 | 8.61 (1H, m), 7.68 (2H, m), 7.20 (1H, m), 3.34 (2H, m), 2.64 (2H, m), 2.5-2.1 (6H, m) and 2.37 (3H, s). |
| 9 | 8.94 (1H, d), 8.59 (1H, t), 8.53 (1H, d), 3.35 (2H, m), 2.65-2.55 (2H, m), 2.4-2.1 (6H, m) and 2.36 (3H, m). |

- 10 8.82 (1H, s), 8.54 (1H, s), 3.37 (2H, m), 2.54 (2H, dd), 2.4-2.1 (6H, m) and 2.35 (3H, s).
- 11 7.78 (1H, d), 7.55 (1H, d), 3.40 (2H, m), 2.70 (2H, m), 2.5-2.1 (6H, m) and 2.37 (3H, s).
- 12 8.49 (1H, d), 7.95 (1H, d), 3.34 (2H, m), 2.4-2.15 (8H, m) and 2.35 (3H, s).
- 13 8.81 (1H, d), 8.56 (1H, dd), 7.84 (1H, dt), 7.31 (1H, dd), 3.72 (3H, s), 3.50 (2H, m), 3.22 (2H, s), 2.5-2.3 (6H, m) and 2.2-2.1 (2H, m).
- 14 8.89 (1H, d), 8.60 (1H, dd), 7.87 (1H, dt), 7.35 (1H, dd), 4.54 (2H, m), 4.50 (2H, s), 3.22 (3H, s) and 2.6-2.25 (8H, m).
- 15 8.59 (1H, d), 7.81 (1H, dd), 7.34 (1H, d), 3.33 (2H, m), 2.4-2.15 (8H, m) and 2.35 (3H, s).
- 16 8.69 (1H, d), 8.51 (1H, d), 7.82 (1H, t), 3.42 (2H, m), 2.4-2.0 (10H, m), 1.5-1.2 (8H, m) and 0.89 (3H, m).
- 17 8.70 (1H, d), 8.52 (1H, d), 7.83 (1H, t), 5.88 (1H, m), 5.18 (2H, m), 3.42 (2H, m), 3.02 (2H, m) and 2.4-2.05 (8H, m).
- 18 8.70 (1H, d), 8.55 (1H, d), 7.80 (1H, t), 3.50 (2H, m), 2.90 (2H, q), 2.5-2.2 (6H, m) and 2.10 (2H, m).
- 19 8.74 (1H, d), 8.61 (1H, d), 8.00 (1H, t), 3.35 (1H, m), 2.4-2.15 (8H, m) and 2.33 (3H, s).
- 20 9.03 (1H, d), 8.82 (1H, d), 8.15 (1H, t), 3.36 (1H, m), 2.4-2.2 (8H, m) and 2.35 (3H, s).
- 21 8.40 (1H, d), 8.21 (1H, d), 7.34 (1H, t), 4.10 (2H, q), 3.31 (2H, m), 2.4-2.15 (8H, m), 2.32 (3H, s) and 1.44 (3H, t).
- 22 8.70 (1H, d), 8.50 (1H, d), 7.80 (1H, t), 3.70 (2H, m), 2.65 (1H, m), 2.35 (2H, m), 2.25 (4H, m), 2.05 (2H, m) and 1.05 (6H, d).
- 23 7.47 (2H, s), 3.32 (2H, m), 2.35-2.15 (8H, m) and 2.32 (3H, s).
- 24 8.39 (1H, d), 7.51 (1H, d), 7.40 (1H, dd), 3.32 (2H, m), 2.4-2.1 (8H, m) and 2.32 (3H, s).
- 25 8.61 (2H, m), 7.49 (2H, m), 3.33 (2H, m), 2.4-2.1 (8H, m) and 2.32 (3H, s).
- 26 8.69 (1H, d), 8.51 (1H, d), 7.81 (1H, t), 4.35 (1H, ddt), 3.41 (2H, m), 3.00 (2H, dt) and 2.4-2.1 (8H, m).
- 27 8.59 (2H, m), 7.98 (1H, d), 7.72 (1H, d), 5.55 (1H, d), 4.36 (1H, m), 4.25 (1H, m), 2.7-2.2 (8H, m).
- 28 8.65 (1H, d), 8.55 (1H, d), 7.75 (1H, t), 4.95 (1H, m), 4.40 (1H, m), 2.65-2.1 (8H, m) and 2.15 (3H, s).
- 29 [in DMSO] 8.79 (1H, brs), 8.70 (1H, d), 8.64 (1H, brs), 8.61 (1H, d), 4.16 (2H, m), 2.7-2.6 (2H, m), 2.45-2.25 (4H, m), and 2.15-2.0 (2H, m).
- 30 8.67 (1H, d), 8.49 (1H, d), 7.80 (1H, t), 3.79 (2H, m), 2.4-2.15 (6H, m), 2.0-1.9 (2H, m) and 1.09 (9H, s).
- 31 9.56 (1H, s), 8.75 (1H, d), 8.55 (1H, d), 7.89 (1H, t), 7.55-7.3 (5H, m), 3.60 (1H, m), 3.38 (1H, m), 2.6-1.9 (8H, m) and 1.54 (3H, s).
- 32 8.72 (1H, d), 8.53 (1H, d), 7.89 (1H, t), 7.53 (2H, m), 7.35-7.2 (3H, m), 6.11 (1H, dd), 5.29 (1H, d), 5.19 (1H, d), 3.63 (1H, m), 3.54 (1H, m), 2.45-1.9 (8H, m) and 1.51 (3H, s).
- 33 8.69 (1H, d), 8.52 (1H, d), 7.82 (1H, t), 7.54 (2H, m), 7.4-7.25 (3H, m), 3.90 (1H, m), 3.70 (1H, d), 3.65 (1H, d), 3.31 (1H, m), 2.5-2.0 (8H, m) and 1.49 (3H, m).
- 34 8.60 (1H, d), 8.50 (1H, d), 7.71 (1H, t), 7.4-7.25 (5H, m), 3.40 (1H, m), 3.32 (1H, m), 2.82 (1H, t), 2.66 (1H, t), 2.3-1.95 (8H, m) and 1.74 (3H, d).
- 35 8.70 (1H, d), 8.54 (1H, d), 7.85 (1H, t), 7.50 (2H, m), 7.4-7.25 (3H, m), 4.31 (1H, d), 4.19 (1H, d), 3.85 (1H, m), 2.34 (1H, m), 2.5-2.15 (8H, m), 2.00 (3H, s) and 1.56 (3H, s).

- 36 8.64 (1H, d), 8.56 (1H, d), 8.20 (1H, s), 7.77 (1H, t), 4.86 (1H, m), 4.32 (1H, m) and 2.6-2.1 (8H, m).
- 37 8.69 (1H, d), 8.52 (1H, d), 7.85 (1H, t), 4.11 (2H, m), 3.65 (2H, hept), 2.55-2.1 (8H, m) and 1.30 (12H, d).
- 38 8.62 (1H, d), 8.52 (1H, d), 7.75 (1H, t), 4.26 (2H, m), 2.5-2.15 (8H, m) and 1.40 (9H, s).
- 39 8.71 (1H, d), 8.53 (1H, d), 7.84 (1H, t), 7.4-7.2 (5H, m), 3.71 (1H, m), 3.49 (1H, q), 3.28 (1H, m), 2.4-2.05 (8H, m) and 1.30 (3H, d).
- 40 8.20 (1H, d), 8.00 (1H, d), 7.11 (1H, t), 3.76 (2H, brs), 3.32 (2H, m), 2.4-2.1 (8H, m) and 2.32 (3H, s).
- 41 8.62 (1H, d), 8.56 (1H, d), 8.22 (1H, t), 7.46 (1H, brs), 3.35 (2H, m), 2.4-2.2 (8H, m), 2.35 (3H, s) and 2.21 (3H, s).
- 42 8.69 (1H, d), 8.54 (1H, d), 7.81 (1H, t), 7.55-7.35 (5H, m), 4.38 (1H, s), 3.94 (1H, m), 3.29 (1H, m) and 2.6-2.1 (8H, m).
- 43 8.69 (1H, d), 8.55 (1H, d), 7.81 (1H, t), 7.45-7.3 (5H, m), 6.91 (1H, m), 5.70 (1H, m), 3.96 (1H, s), 3.60 (1H, m), 3.35 (1H, m), 2.5-2.2 (6H, m) and 2.05-1.9 (2H, m).
- 44 8.77 (2H, m), 8.16 (1H, t), 3.32 (2H, m), 2.4-2.1 (8H, m) and 2.32 (3H, s).
- 45 9.02 (1H, d), 8.83 (1H, m), 8.07 (1H, m), 3.38 (2H, m) and 2.4-2.05 (11H, m).
- 46 [in DMSO] 8.72 and 8.62 (1H, m), 8.58 (1H, m), 8.14 and 7.86 (1H, m), 5.39 (1H, m), 4.20 (1H, m) 2.7-2.0 (8H, m).
- 47 8.65 (1H, d), 8.58 (1H, d), 7.78 (1H, t), 4.51 (2H, m), 2.6-2.2 (8H, m).
- 48 8.69 (1H, d), 8.52 (1H, d), 7.81 (1H, t), 5.85 (1H, t), 3.45 (1H, m), 2.71 (2H, dt) and 2.5-2.0 (8H, m).
- 49 8.69 (1H, d), 8.52 (1H, d), 7.82 (1H, t), 7.35-7.15 (5H, m), 3.45 (2H, m), 2.79 (2H, m), 2.61 (2H, m) and 2.4-2.05 (8H, m).
- 50 8.39 (1H, m), 8.05 (1H, m), 7.59 (1H, m), 3.43 (2H, m) 2.5-2.1 (8H, m) and 2.40 (3H, s).
- 51 8.70 (1H, d), 8.52 (1H, d), 7.83 (1H, t), 7.4-7.2 (5H, m), 3.57 (2H, s), 3.40 (2H, m) and 2.45-2.2 (8H, m).
- 52 8.63 (1H, d), 8.50 (1H, d), 7.77 (1H, t), 3.60 (2H, m), 3.42 (2H, m) and 2.5-2.2 (8H, m).
- 53 8.71 (1H, d), 8.53 (1H, d), 8.10 (1H, t), 7.90 (2H, m), 7.39 (2H, m), 3.45 (2H, m), 3.30 (2H, m) and 2.4-2.2 (8H, m).
- 54 8.69 (1H, d), 8.53 (1H, d), 7.83 (1H, t), 7.42 (1H, dd), 7.22 (1H, m), 7.09 (1H, t), 3.50 (2H, s), 3.36 (2H, m) and 2.45-2.15 (8H, m).
- 55 8.71 (1H, d), 8.52 (2H, m), 7.86 (1H, t), 7.69 (1H, dt), 7.54 (1H, d), 7.19 (1H, m), 3.73 (2H, s), 3.43 (2H, m) and 2.5-2.2 (8H, m).
- 56 8.70 (1H, d), 8.52 (1H, d), 7.85 (1H, t), 7.00 (1H, s), 3.66 (2H, s), 3.49 (2H, m), 2.70 (3H, s) and 2.45-2.15 (8H, m).
- 57 8.61 (1H, d), 8.52 (1H, d), 7.78 (1H, t), 3.33 (2H, m), 3.29 (2H, s), 2.5-2.1 (8H, m), 2.38 (3H, s) and 2.32 (3H, s).
- 58 8.70 (1H, d), 8.52 (1H, d), 7.83 (1H, t), 6.74 (1H, d), 6.65 (1H, d), 3.65 (2H, s), 3.46 (2H, m) and 2.45-2.1 (8H, m).
- 59 8.65 (1H, d), 8.50 (1H, d), 7.80 (1H, t), 3.94 (2H, s), 3.55 (2H, m) and 2.5-2.25 (8H, m).
- 60 8.65 (1H, d), 8.41 (1H, d), 7.60 (1H, dt), 3.44 (2H, m), 2.4-2.15 (8H, m) and 2.35 (3H, s).
- 61 8.69 (2H, m), 7.84 (1H, t), 7.10 (2H, m), 6.40 (2H, m), 3.34 (2H, m), 2.4-2.15 (8H, m) and 2.33 (3H, s).

- 62 8.79 (1H, d), 8.75 (1H, d), 8.05 (1H, m), 4.74 (1H, d), 4.31 (1H, d), 3.94 (2H, q), 3.40 (2H, m), 2.5-2.2 (11H, m) and 1.44 (3H, t).
- 63 9.09 (1H, d), 8.99 (1H, d), 8.39 (1H, t), 3.37 (2H, m), 2.69 (3H, s), 2.45-2.15 (8H, m) and 2.34 (3H, s).
- 64 8.76 (1H, d), 8.62 (1H, d), 7.94 (1H, m), 3.34 (2H, m), 3.24 (1H, s), 2.45-2.15 (8H, m) and 2.35 (3H, s).
- 65 8.80 and 8.70 (1H, m), 8.29 (1H, m), 7.95 (1H, m), 7.71 (1H, m), 7.56 (1H, m), 7.4-7.1 (3H, m), 5.65 and 5.29 (1H, m), 4.91 and 4.75 (1H, m), 4.30 and 4.11 (1H, m), 3.30 (1H, m), 2.8-2.0 (8H, m) and 1.73 and 1.62 (3H, m).
- 66 8.61 (1H, d), 8.52 (1H, d), 7.79 (1H, t), 7.5-7.15 (5H, m), 3.01 (1H, m), 2.91 (1H, m), 2.79 (1H, m), 2.39 (2H, m), 2.3-1.6 (12H, m), 1.09 (3H, d) and 0.91 (3H, d).
- 67 8.69 (1H, d), 8.52 (1H, d), 7.81 (1H, t), 3.55 (2H, m), 3.35 (2H, s) and 2.5-2.1 (8H, m).
- 68 8.69 (1H, d), 8.52 (1H, d), 7.81 (1H, t), 4.20 (2H, m), 3.61 (1H, m), 3.49 (1H, m), 3.23 (1H, q), 2.45-2.0 (8H, m), 1.31 (3H, d) and 1.29 (3H, t).
- 69 8.50 (1H, d), 8.35 (1H, d), 7.7-7.5 (2H, m), 6.40 (2H, dd), 6.25 (1H, dd), 4.70 (2H, m), 2.6-2.5 (2H, m) 2.45-2.20 (6H, m).
- 70 8.51 (1H, d), 8.49 (1H, d), 7.70 (1H, t), 4.45 (2H, m), 2.6-2.5 (4H, m) and 2.4-2.2 (4H, m).
- 71 8.70 (1H, d), 8.55 (1H, d), 7.85 (1H, t), 4.25 (2H, m) and 2.6-2.2 (8H, m).
- 72 8.29 (1H, d), 8.10 (1H, d), 7.90 (1H, m), 4.15 (2H, m) 2.6-2.25 (8H, m).
- 73 8.75, 8.65 and 8.50 (1H, m), 7.90 and 7.80 (1H, m), 3.80 and 3.70 (2H, m), 3.6 and 3.5 (3H, m) and 2.7-1.8 (8H, m).
- 74 8.65 (1H, d), 8.51 (1H, d), 7.80 (1H, t), 4.16 (2H, q), 3.41 (2H, m), 2.69 (2H, t), 2.48 (2H, t), 2.4-2.05 (8H, m) and 1.27 (3H, t).
- 75 8.76 (1H, m), 8.52 (1H, m), 7.98 (1H, m), 3.85 (2H, m), 3.01 (2H, t), 2.8-2.2 (10H, m).
- 76 8.64 (1H, d), 8.51 (1H, d), 7.80 (1H, t), 4.18 (4H, m), 3.60 (2H, m), 2.78 (2H, d), 2.45-2.05 (8H, m) and 1.36 (6H, m).
- 77 [in DMSO] 8.89 (1H, d), 8.62 (1H, d), 8.29 (1H, t), 4.50 (2H, m), 3.48 (2H, d) and 2.85-2.45 (8H, m).
- 78 8.69 (1H, d), 8.51 (1H, d), 7.80 (1H, t), 3.46 (2H, m), 2.64 (2H, m), 2.51 (2H, m) and 2.45-2.0 (8H, m).
- 79 9.65 (1H, d), 8.51 (1H, d), 7.79 (1H, t), 3.91 (2H, m), 2.45-2.2 (8H, m) and 1.28 (6H, s).
- 80 8.55 (1H, d), 8.3 (1H, d) 7.45 (1H, t), 3.35 (2H, m), 2.4-2.1 (8H, m) and 2.3 (3H, s).
- 81 8.69 (1H, d), 8.52 (1H, d), 7.81 (1H, t), 3.85 (1H, m), 3.60 (1H, m), 3.50 (1H, q), 2.5-2.1 (8H, m) and 1.51 (3H, d).
- 82 8.81 (1H, d), 8.71 (1H, d), 7.92 (1H, t), 7.6-7.35 (5H, m), 7.25 (1H, dd), 4.80 (1H, dd), 4.61 (2H, m), 4.49 (1H, dd) and 2.55-2.15 (8H, m).
- 83 8.79 (2H, m), 7.99 (1H, t), 7.6-7.35 (5H, m), 3.75 (2H, m), 2.55-2.2 (6H, m) and 2.05-1.85 (2H, m).
- 84 8.55 (1H, m), 8.40 (1H, m), 7.55 (1H, m), 7.23 (1H, dd), 4.80 (1H, dd), 4.59 (1H, m), 4.49 (1H, dd), 2.6-2.1 (8H, m) and 2.39 (3H, s).
- 85 8.64 (1H, d), 8.52 (1H, d), 7.70 (1H, t), 3.91 (1H, m), 2.55-2.15 (8H, m) and 1.52 (6H, s).
- 86 9.19 (1H, d), 8.90 (1H, d), 8.31 (1H, t), 7.24 (1H, dd), 4.82 (1H, dd), 4.63 (2H, m), 4.51 (1H, dd), 4.44 (2H, q), 2.6-2.2 (8H, m) and 1.43 (3H, t).

EXAMPLE 87

This Example illustrates an emulsifiable concentrate composition which is readily convertible by dilution with water into a liquid preparation suitable for spraying purposes. The concentrate has the following composition:

	% Weight
Compound No.1	25.5
SYNPERONIC NP13	2.5
Calcium dodecylbenzenesulphonate	2.5
10 AROMASOL H	70

EXAMPLE 88

This Example illustrates an emulsifiable concentrate composition which is readily convertible by dilution with water into a liquid preparation suitable for spraying purposes. The concentrate has the following composition:

	% Weight
Compound No.5	50.0
SYNPERONIC NP13	6.0
Calcium dodecylbenzenesulphonate	4.0
20 AROMASOL H	40.0

EXAMPLE 89

This Example illustrates an emulsifiable concentrate composition which is readily convertible by dilution with water into a liquid preparation suitable for spraying purposes. The concentrate has the following composition:

	% Weight
25 Compound No.9	1.0
SYNPERONIC OP10	3.0
Calcium dodecylbenzenesulphonate	2.0
AROMASOL H	94.0

EXAMPLE 90

30 This Example illustrates a wettable powder composition which is readily convertible by dilution with water into a liquid preparation suitable for spraying purposes. The wettable powder has the following composition:

- 62 -

	% Weight
Compound No.13	25.0
Silica	25.0
Sodium lignosulphonate	5.0
5 Sodium lauryl sulphate	2.0
Kaolinite	43.0

EXAMPLE 91

This Example illustrates a wettable powder composition which is readily convertible by dilution with water into a liquid preparation suitable for spraying purposes. The wettable powder has the following composition:

	% Weight
Compound No.17	1.0
Sodium lignosulphonate	5.0
Sodium lauryl sulphate	2.0
15 Kaolinite	92.0

EXAMPLE 92

This Example illustrates a wettable powder composition which is readily convertible by dilution with water into a liquid preparation suitable for spraying purposes. The wettable power has the following composition:

	% Weight
Compound No.21	40.0
Silica	40.0
Calcium lignosulphonate	5.0
25 Sodium lauryl sulphate	2.0
Kaolinite	13.0

EXAMPLE 93

This Example illustrates a dusting powder which may be applied directly to plants or other surfaces and comprises 1% by weight of Compound No.25 and 99% by weight of talc.

EXAMPLE 94

This Example illustrates a concentrated liquid formulation suitable for application by ultra low volume techniques after mixing with paraffinic diluents.

	% Weight
5 Compound No.29	90.0
SOLVESSO 200	10.0

EXAMPLE 95

This Example illustrates a concentrated liquid formulation suitable for application by ultra low volume techniques after mixing with paraffinic diluents.

10	% Weight
Compound No.33	25.0
SOLVESSO 200	75.0

EXAMPLE 96

15 This Example illustrates a concentrated liquid formulation suitable for application by ultra low volume techniques after mixing with paraffinic diluents.

	% Weight
Compound No.37	10.0
SOLVESSO 200	90.0

EXAMPLE 97

20 This Example illustrates a liquid formulation suitable for application (undiluted) by ultra low volume techniques.

	% Weight
Compound No.41	15
25 Cotton seed oil	50
SOLVESSO 200	35

EXAMPLE 98

This Example illustrates a capsule suspension concentrate which is readily convertible by dilution with water to form a preparation suitable for application as an aqueous spray.

30	% Weight
Compound No.45	10.0

- 64 -

	Alkylbenzene solvent (e.g. AROMASOL H)	5.0
	Toluene di-isocyanate	3.0
	Ethylenediamine	2.0
	Polyvinyl alcohol	2.0
5	Bentonite	1.5
	Polysaccharide (e.g. KELTROL)	0.1
	Water	76.4

EXAMPLE 99

This Example illustrates a capsule suspension concentrate which is readily
 10 convertible by dilution with water to form a preparation suitable for application as an aqueous spray.

		% Weight
	Compound No.49	1.0
	Alkylbenzene solvent (e.g. AROMASOL H)	10.0
15	Toluene di-isocyanate	3.0
	Ethylenediamine	2.0
	Polyvinyl alcohol	2.0
	Bentonite	1.5
	Polysaccharide (e.g. KELTROL)	0.1
20	Water	80.4

EXAMPLE 100

A ready for use granular formulation:

		% Weight
	Compound No.4	0.5
25	SOLVESSO 200	0.2
	nonylphenol ethoxylate (eg Synperonic NP8)	0.1
	Calcium carbonate granules (0.3-0.7 mm)	99.2

30 EXAMPLE 101

An aqueous suspension concentrate:

% Weight

- 65 -

	Compound No.8	5.0
	Kaolinite	15.0
	Sodium lignosulphonate	3.0
	nonylphenolethoxylate	1.5
5	(eg Synperonic NP 8)	
	propylene glycol	10.0
	Bentonite	2.0
	Polysaccharide (eg Keltrol)	0.1
	Bactericide (eg Proxel; Proxel	0.1
10	is a registered Trade Mark)	
	Water	63.3

EXAMPLE 102

A ready for use dust (D.P.) made from a concentrate

Concentrate:

15		% Weight
	Compound No.12	10
	Silica	20
	Magnesium Carbonate	70

20 Dust Example containing 1% active ingredient:

	Above concentrate	10
	Talc	90

EXAMPLE 103

This Example illustrates a ready for use granule formulaton.

25		% Weight
	Compound No.16	5
	Synperonic NP8	2
	Pumice granules (20/40 BS Mesh)	93

30 EXAMPLE 104

This Example illustrates a water dispersible granule formulation.

% Weight

- 66 -

Compound No.20	5
Silica	5
Sodium lignosulphate	10
Sodium dioctylsulphosuccinate	5
5 Sodium acetate	10
Montmorillonite powder	65

EXAMPLE 105

This Example illustrates the insecticidal properties of the compounds of Formula I. The activity of the the compounds of Formula I was determined using a variety of pests. The pests
 10 were treated with a liquid composition containing 500 parts per million (ppm) by weight of the compound unless otherwise stated. The compositions were made by dissolving the compound in acetone and ethanol (50:50) mixture and diluting the solutions with water containing 0.05% by weight of a wetting agent sold under the trade name "SYNPERONIC" NP8 until the liquid composition contained the required concentration of the compound. "SYNPERONIC" is a
 15 Registered Trade Mark.

The test procedure adopted with regard to each pest was basically the same and comprised supporting a number of the pests on a medium which was usually a substrate, a host plant or a foodstuff on which the pests feed, and treating either or both the medium and the pests with the compositions. The mortality of the pests was then assessed at periods usually varying
 20 from two to five days after the treatment.

The results of the tests against peach aphid (*Myzus persicae*) are presented in Table II. The results indicate a grading of mortality (score) designated as A, B or C wherein C indicates less than 40% mortality, B indicates 40-79% mortality and A indicates 80-100% mortality; "-" indicates that either the compound was not tested or no meaningful result was obtained. In this
 25 test Chinese cabbage leaves were infested with aphids, the infested leaves were sprayed with the test composition, and the mortality assessed after 3 days.

Information regarding the pest species, the support medium or food, and the type and duration of the test is given in Table III. The pest species is designated by a letter code.

TABLE II

Comp'd No	Score	Comp'd No	Score	Comp'd No	Score	Comp'd No	Score
1	A	2	A	3	A	4	A
5	A	6	C	7	C	8	A
9	A	10	A	11	A	12	A
13	B	14	A	15	A	16	C
17	C	18	A	19	C	20	C
21	A	22	A	23	A	24	A
25	A	26	B	27	A	28	C
29	B	30	A	31	C	32	A
33	A	34	A	35	A	36	A
37	A	38	A	39	C	40	A
41	A	42	A	43	A	44	A
45	C	46	A	47	A	48	A
49	A	50	A	51	A	52	C
53	A	54	A	55	A	56	A
57	A	58	C	59	A	60	A
61	A	62	A	63	C	64	A
65	C	66	C	67	A	68	A
69	A	70	A	71	A	72	A
73	C	74	A	75	A	76	A
77	A	78	A	79	A	80	A
81	A	82	A	83	A	84	A
85	A	86	A	87	A	88	A
89	C	90	A	91	A	92	A
93	A	94	C	95	A	96	A
97	C	98	A	99	A	100	A
101	A	102	A	103	A	104	A
105	B	106	A	107	A	108	A

- 68 -

109	A	110	A	111	A	112	B
113	A	114	A	115	A	116	A
117	A	118	A	119	A	120	A
121	A	122	A	123	A	124	A
125	A	126	C	127	-	128	A
129	A	130	A	131	A	132	A
133	A	134	A	135	A	136	C
137	C	138	A	139	A	140	A
141	A	142	A	143	A	144	A
145	A	146	A	147	A	148	B
149	A	150	A	151	A	152	A
153	A	154	B	155	A	156	A
157	C	158	A	159	A	160	A
161	A	162	C	163	C	164	A
165	A	166	A	167	A	168	A
169	A	170	B	171	A	172	A
173	A	174	A	175	A	176	A
177	A	178	A	179	A	180	A
181	A	182	A	183	A	184	A
185	A	186	A	187	C	188	A
189	A	190	-	191	A	192	A
193	A	194	A	195	A	196	A
197	A	198	A	199	A	200	A
201	A	202	A	203	A	204	A
205	A	206	A	207	A	208	A
209	A	210	A	211	A	212	A
213	A	214	A	215	A	216	-
217	A	218	A	219	-	220	A
221	A	222	C	223	C	224	-
225	A	226	-	227	A	228	A
229	A	230	A	231	A	232	A
233	A	234	A	235	A	236	A

- 69 -

237	A	238	A	239	A	240	A
241	A	242	A	243	A	245	A
246	A	247	A	248	A	249	A
250	A	251	A	252	A	253	A
254	A	255	A	256	A	257	A
258	A	259	A	260	A	261	A
262	A	263	A	264	A	265	A
266	A	267	A	268	A	269	A
270	A	271	A	272	A	273	A
274	A	275	A	276	A	277	A
278	A	279	A	280	A	281	A
282	A	283	A	284	A	285	A
286	A	287	A	288	A	289	A
290	A	291	-	292	A	293	A
294	A	295	-	296	A	297	A
298	A	299	A	300	A	301	A
302	A	303	-	304	A	305	-
306	-	307	-	308	-	309	A
310	A	311	A	312	A	313	A
314	A	315	A	316	A	317	A
318	A	319	A	320	A	321	A

In tests against tobacco budworm (Heliothis virescens, larvae) the following compounds scored A or B.

Compounds 2, 8, 14, 18, 23, 66, 72, 95, 99, 102, 104, 120, 131, 156, 169, 170, 227, 229, 231,
 5 234, 236, 243, 260, 262, 270, 274, 312.

In tests against root knot nematodes (Meloidogyne incognita) the following compounds scored A or B.

Compounds 36, 55, 71, 77, 94, 99, 120, 160, 207, 233, 237, 238, 253, 257, 271, 312, 317, 318.











In tests against red spider mite (Tetranychus urticae) the following compounds scored A or B.

10 Compounds 12, 13, 22, 23, 25, 34, 37, 39, 47, 53, 63, 64, 66, 87, 90, 99, 101, 106, 120, 135,
 142, 186, 193, 195, 199, 201, 207, 208, 236, 237, 239, 245, 247, 249, 280, 283, 310 to 321.

In tests against Whitefly (Bemesia tabaci) the following compounds were particularly effective.


Compounds 33, 34, 36, 56, 64, 68, 69, 70, 72, 74, 76, 77, 81, 90, 93, 99, 227 to 275.

The chemical formulae referred to in the preceding description are set out below.




CLAIMS

1. A compound of formula (I):



wherein R¹ represents a group of formula (A)




where each of W, X, Y and Z and Z

represents either a group CR or the nitrogen atom, provided that not more than two of W, X, Y and Z represent the nitrogen atom and where each R present is independently selected from hydrogen and halogen atoms and cyano, amino, hydrazino, acylamino, hydroxy, alkyl, hydroxyalkyl, alkoxy, haloalkyl, haloalkoxy, alkenyl, alkenyloxy, alkoxyalkenyl, alkynyl, carboxylic acyl, alkoxycarbonyl, aryl and heterocyclyl groups, said groups comprising up to 6 carbon atoms, and wherein R² represents hydrogen or cyano or a group selected from alkyl, aryl, heteroaryl, aralkyl, heteroarylalkyl, alkenyl, aralkenyl, alkynyl, alkoxycarbonyl, alkanesulfonyl, arenesulfonyl, alkanyloxycarbonyl, aralkyloxycarbonyl, aryloxycarbonyl, heterocyclylalkyl, carbamyl or dithiocarboxyl groups, said groups comprising from 1 to 15 carbon atoms, said groups being optionally substituted with one or more substituents selected from, halogen, cyano, carboxyl, carboxylic acyl, carbamyl, alkoxycarbonyl, alkoxy, alkylenedioxy, hydroxy, nitro, haloalkyl, alkyl, amino, acylamino, imidate and phosphonato groups; and acid addition salts, quaternary ammonium salts and N-oxides derived therefrom.


2. A compound according to claim 1 wherein R¹ represents a halo-substituted phenyl, pyridyl or diazinyl group.
3. A compound according to claim 1 where R¹ represents an optionally halogen substituted phenyl group or an optionally halogen substituted pyridyl, pyridazinyl or pyrazinyl group and R² represents hydrogen or a C₁₋₆ alkyl, alkenyl, alkynyl, phenyl, benzyl, pyridylmethyl, thienylmethyl, thiazolylmethyl group which may be optionally substituted with one or more alkyl, alkoxy, alkoxycarbonyl, cyano, optionally substituted alkane sulphonyl groups or halogen atoms; and acid addition salts thereof.
4. A compound according to claim 1 wherein R¹ is a halo-substituted pyridyl group.
5. A compound according to claim 1 wherein R² is a hydrogen or haloalkyl, haloalkenyl or haloaralkyl group.
6. A compound according to claim 4 wherein R² is a haloalkyl or haloalkenyl group.
7. A compound according to claim 6 wherein R² is a fluoroalkyl or fluoroalkenyl group.
8. A compound according to claim 4 where R¹ is a 5-halopyrid-3-yl group.
9. A compound according to claim 8 wherein R² is fluoroethyl, difluoroethyl or trifluoroethyl.
10. An insecticidal acaricidal and nematicidal composition comprising an insecticidally, acaricidally or nematicidally effective amount of a compound according to claim 1.
11. A method of combating and controlling acarine or nematode pests at a locus which comprises treating the pests or the locus of the pests with an effective amount of a composition according to claim 10.
12. A method according to claim 11 wherein the pests are insect pests of growing plants.

13. A method of preparing a compound of formula (I) where R^2 is not hydrogen which comprises reacting a compound of formula (II):




with a compound of formula R^2L where L is a leaving group in the presence of a base.

14. A process according to claim 13 wherein L represents halide or triflate.
- 10
15. A process of preparing a compound of formula (I) which comprises reacting a compound of formula (VI):




with a compound of formula R^1Hal where Hal is a halide in the presence of a base.

- 15
16. A process of preparing a compound of formula (VI):




which comprises reacting a compound of formula (VIII):



with a compound of formula R^2L where L is a leaving group in the presence of a base.

5

17. A compound of formula (VI):



wherein R^2 has any of the meanings given in claim 1 provided that R^2 is not methyl, benzyl or trichloroethyl.

10

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 96/01151

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D451/02 A01N43/40

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07D A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JOURNAL OF MEDICINAL CHEMISTRY, vol. 18, no. 5, 1975, WASHINGTON US, pages 496-501, XP002008338 S.J.DAUM ET AL.: "Analgesic activity of the epimeric tropane analogs of meperidine." * page 497, 500 : compounds 7 and 8 and their HCl salts * -----	1

☐ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

12 July 1996

Date of mailing of the international search report

25.07.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Van Bijlen, H